



# *The MoHole*

## A Crustal Journey and Mantle Quest

### *Workshop Report*

Kanazawa, Japan, 3-5 June 2010



#### **Workshop Steering Committee :**

Benoît Ildefonse, Natsue Abe, Yoshio Isozaki (Chairs)  
Donna K. Blackman, J. Pablo Canales, Shuichi Kodaira, Greg Myers, Mladen R. Nedimovic, Damon A.H. Teagle, Susumu Umino (host), Douglas S. Wilson

#### **Report Writing Committee :**

B. Ildefonse (Chair), N. Abe, D.K. Blackman, J.P. Canales, Y. Isozaki, S. Kodaira, G. Myers, K. Nakamura, M. Nedimovic, N. Seama, A. Skinner, E. Takazawa, D.A.H. Teagle, M. Tominaga, S. Umino, D.S. Wilson, M. Yamao

# Contents

<b>Executive Summary.....</b>	<b>1</b>
<b>1. Introduction.....</b>	<b>4</b>
<b>2. Scientific Rationale for the MoHole, a Crustal Journey and Mantle Quest.....</b>	<b>5</b>
2.1. The Mohorovičić Discontinuity.....	5
2.2. Formation of the lower crust.....	6
2.3. Composition and hydration of the ocean crust.....	7
2.4. Limits and controlling factors of life.....	8
2.5. Physical and chemical nature of the upper mantle.....	8
<b>3. Geophysical Site Surveys.....</b>	<b>10</b>
3.1. Scientific Requirements.....	10
3.2. Technological Constraints.....	10
3.3. Potential Sites.....	12
3.3.1. Cocos Plate (aka 1256 area).....	14
3.3.2. Off Southern/Baja California.....	15
3.3.3. Hawaii.....	16
3.4. Large Scale Surveys: Finding the Right Project Area.....	17
3.5. Small-Scale Surveys: Detailed Imaging of Project Area and Target.....	18
<b>4. Technology &amp; Engineering Development.....</b>	<b>19</b>
4.1. Mud Circulation Technology.....	19
4.2. Logging/ Coring.....	20
4.3. Drill Bit, Drill String, and Core Barrels.....	21
4.4. Drilling Mud.....	21
4.5. Casing/Cementing.....	22
4.6. Assessing the Past Experience in Crustal Drilling.....	23
4.7. Accumulating Experience in Hard-Rock Drilling.....	23
<b>5. Operations.....</b>	<b>24</b>
5.1. Professional Planning.....	24
5.2. Site Characterization Data.....	24
5.3. Well Design Considerations.....	25
5.3.1. Pilot Hole Options.....	25
5.3.2. Well Control and Borehole Pressure Management.....	25
5.3.3. Mud program.....	25
5.3.4. Casing design.....	25
5.3.5. Cementing plan.....	26
5.3.6. Directional drilling options.....	26
5.4. Downhole drilling tool selection.....	26
5.5. Downhole Measurements.....	26
5.5.1. Drill collar measurements - Logging While Drilling/Measurement While Drilling.....	26
5.5.2. Wireline tools.....	27
5.5.3. VSP experiment.....	27
5.5.4. Long-term monitoring.....	27
5.6. Logistics.....	27
<b>6. Keys for success.....</b>	<b>28</b>
6.1. Geophysical surveys and Hole location.....	28
6.2. Scientific coring, sampling and measurements.....	28
6.3. Broadening the scientific spectrum.....	30



6.4. Drilling technology .....	30
6.5. Industrial engagement.....	31
6.6. Public engagement, outreach, education.....	31
6.7. Anticipated time table for the MoHole project.....	32
<b>7. References .....</b>	<b>33</b>
<b>8. Workshop Agenda.....</b>	<b>35</b>
Thursday, June 3 - Day 1.....	35
Friday, June 4 - Day 2 .....	36
Saturday, June 5 - Day 3.....	36
<b>9. Workshop Participants .....</b>	<b>37</b>
<b>Appendices .....</b>	<b>39</b>
Appendix 1 - Scientific Objectives of Mission Moho proposal (2007).....	40
1. Primary Motivation for Mission Moho.....	41
2. Road to the Moho.....	42
3. Scientific Objectives.....	43
References.....	52
Appendix 2 - Langseth Workshop White Paper (2010) .....	58
Appendix 3 - INVEST Meeting White Papers.....	61
Appendix 4 - Links to Earlier Scientific Planning Documents.....	63
Appendix 5 - Workshop Presentation slides .....	64
<i>Introduction, scientific rationale for the MoHole</i> .....	64
<i>Available Site survey data in the Pacific</i> .....	64
<i>Current understanding of technological feasibility</i> .....	64
<i>Drilling and coring at high temperatures</i> .....	64

## Executive Summary

Drilling an ultra-deep hole in an intact portion of oceanic lithosphere, through the crust to the Mohorovičić discontinuity (the 'Moho'), and into the uppermost mantle is a long-standing ambition of scientific ocean drilling, and remains essential to answer fundamental questions about the dynamics of the Earth and global elemental cycles. The 2010 MoHole workshop had two interconnected objectives, which have been discussed jointly between ocean lithosphere specialists, marine geophysicists, and engineers:

- to initiate a roadmap for technology development, and the project implementation plan, to achieve the deep drilling objectives of the MoHole project,
- to identify potential MoHole sites in Pacific fast-spread crust, where the scientific community will focus geophysical site survey efforts over the next few years.

New deep drilling technology now make it possible to fulfill our long term aspiration to drill completely through intact oceanic crust and a significant distance into the upper mantle, and address a number of first-order scientific goals: what is the geological nature of the Moho? How is the oceanic crust formed at mid-ocean ridges, and what processes influence its subsequent evolution? What are the geophysical signatures of these magmatic, tectonic, hydrothermal, and biogeochemical chemical processes? What is the global composition of the oceanic crust, and what are the magnitudes of interactions with the oceans and biosphere, and their influence on global chemical cycles? What are the limits of life, the factors controlling these limits, and the changes in biological community composition with depth? What is the physical and chemical nature of the uppermost mantle, and how does it relate to the overlying magmatic crust?

The selected MoHole target would ideally meet a suite of scientific requirements including fast spreading rate, simple tectonic setting, "normal" crustal seismic structure, and strong reflectivity of Moho. Several technological constraints limit the range or possible sites, including in particular the trade-off between seafloor depth, which should be small enough to allow using mud re-circulating technologies, and temperature at Moho/upper mantle depths, which should be low enough ( $\sim \leq 250^\circ\text{C}$ ) to allow ultra deep drilling (6000 to 7000 meters below seafloor) in basement. The workshop participants discussed three areas in the Pacific Basin:

- The Cocos plate region encompasses a section of the Cocos Plate off Central America with crustal ages between 15 and 25 Ma. This area includes ODP Hole 1256D, a site of on-going deep drilling into intact ocean crust. This region sits within a corridor that includes a complete tectonic plate life cycle, making it an excellent candidate for understanding ocean crust evolution from a spreading center to subduction. However, temperature at Moho depth at Site 1256 may exceed technology limitations.
- The Off Southern/Baja California region encompasses a section of the eastern Pacific Plate between  $\sim 10\text{--}33^\circ\text{N}$ , and  $\sim 130\text{--}115^\circ\text{W}$ , with crustal ages 20-35 Ma. Very little modern geophysical information exists in this region.
- The Hawaii region is located north of Oahu in the flexural arch, where the crust is 80 Ma, and was formed at an intermediate half spreading rate of 35-40 mm/yr. This site offers the lowest temperature at Moho depth, but crustal structure is potentially

affected by hotspot volcanism, and its cretaceous age makes it difficult to relate geochemical changes to modern ocean chemistry or conditions.

Short-term priority is to conduct large-scale geophysical surveys (spatially coincident MCS, wide-angle OBS, multi-beam bathymetry, gravity, heat flow, and magnetic anomaly data) in the three selected regions, which will lead to the identification of a MoHole target that best satisfies the listed requirements. The first large-scale survey will be in the Off Southern/Baja California region; baseline reconnaissance seismic data are urgently required to assess whether this area can possibly meet the requirements. After an appropriate drilling target has been identified, the community should conduct detailed seismic surveys in its vicinity, to accurately image intracrustal reflectors and Moho, and to assess crustal structure and thickness variability and upper mantle velocity structure/anisotropy. Although JAMSTEC is taking the lead for conducting the first large-scale survey in 2011, other partners should make efforts to conduct or contribute to other large- and small-scale surveys. The scope and costs of these are too large to be undertaken by a single nation or funding agency; international collaboration is required.

Engineering efforts must be directed to ensure that our scientific goals are achieved. The technology and engineering development should be launched as soon as possible in conjunction with site-survey activities. To do so, establishing a realistic roadmap of technological development and testing is imperative. To drill an ultra deep borehole, the provision for continuous mud circulation is a top priority technology requirement. Various options were discussed at the workshop, and will need to be fully evaluated. Other major areas requiring engineering consideration include logging and coring in high temperature environment, drill bits (specifically designed for abrasive, hard rocks) and drill string (high tensile strength), drilling mud (developed for high temperature environment), and casing/cementing materials and strategies (specifically designed, ideally to the bottom of the hole).

Major challenges of the MoHole project will be associated with collecting the cored material, making in-situ measurements, installing casing and keeping the borehole open for successive episodes of deepening in a multiyear, multiphase operation. To gear up for operations, all issues related to drilling, casing, coring, and logging must be adequately explored and included in a comprehensive and complete operation plan, as soon as the site characteristics are known. The well design of the primary site may require data from a pilot hole, to properly evaluate parameters such as mud weights and casing set points. The pilot hole may be either a separate hole, or simply a pilot section of the main hole.

The keys for a successful MoHole project include scientific considerations, as well as technology development, industry engagement, and public engagement through outreach activities and education. The MoHole project will be one of the largest scientific endeavors in Earth science history, and this formidable challenge should provide precious opportunities to diverse scientific and technology communities. It will be essential to share these opportunities and achievements across a broad spectrum of Earth and Life scientists. Many of our primary scientific goals will require continuous core samples. To be regarded as successful, the MoHole project must at least return:

- Continuous cores of critical intervals (lithological boundaries; the region identified by seismic imaging as the Moho; the lower ~500m of the mafic and ultramafic

cumulate rocks in the oceanic crust; ~500m of peridotites and associated lithologies in the uppermost mantle below the Moho; sufficient sections of the igneous oceanic crust to test models of crustal accretion, melt movement, resolve the geometry and intensity of hydrothermal circulation, and document the limits and activity of the deep microbial biosphere).

- A continuous, comprehensive suite of geophysical logs and borehole experiments to measure in situ physical properties, to acquire borehole images, and to identify key geophysical and lithologic regions and transitions throughout the ocean crust and into the upper mantle.

Measurements of temperature and chemical compositions of the fluids are required together with biological analyses. Ultimately, it will be essential to integrate core/log/survey data in a comprehensive synthesis study of the Project Area. Post-drilling studies, such as successive fluid measurements/sampling, in situ microbial incubation in the borehole, or VSP experiments should also be performed.

As the oil and gas industry conducts operations in increasingly deep water, continuous collaboration with industry, and introduction of new technologies to the MoHole project will be keys to success. It will be necessary to establish a strategy to engage the industry in the project, exchange personnel, and plan joint development work.

Another key component of the success of a Mohole project will be to improve public support and understanding of the scientific goals and excitement of the project. Engaging the public through outreach and education activities, as well as being pro-active in advertising the project to the wider scientific community, will be essential. One tool to greatly increase the public and scientific profile of the project should be a dedicated, multi-language MoHole website, that should be implemented as soon as formal scoping activities commence.

The size and duration (10 years or more) of the MoHole project will require an appropriately funded, centralized science operations and engineering management group to oversee the project from start to successful completion. This project office will be vital key to success, and in order to succeed, it must be a truly international effort staffed by the best available project managers, engineers, and scientists.

## 1. Introduction

The global system of mid-ocean ridges and the new oceanic lithosphere formed at these spreading centres are the principal pathways for energy and mass exchange between the Earth's interior, hydrosphere, and biosphere. Bio-geochemical reactions between the oceans and oceanic crust continue from ridge to subduction zone, and the physical and chemical changes to the ocean lithosphere provide inventories of these thermal, chemical and biological exchanges. Drilling an ultra-deep hole in an intact portion of oceanic lithosphere, through the crust to the Mohorovičić discontinuity (the 'Moho'), and into the uppermost mantle is a long-standing ambition of scientific ocean drilling (e.g., Bascom, 1961; Shor, 1985; Ildefonse et al., 2007), and remains essential to answer fundamental questions about the dynamics of the Earth and global elemental cycles.

The 2010 MoHole workshop in Kanazawa, Japan, followed on from several recent scientific planning meetings on ocean lithosphere drilling, in particular the Mission Moho Workshop in 2006 (<http://www.iodp.org/mission-moho-workshop>; Christie et al., 2006; Ildefonse et al., 2007), and the "Melting, Magma, Fluids and Life" meeting in 2009 (<http://www.interridge.org/WG/DeepEarthSampling/workshop2009>, Teagle et al., 2009). Participants to these previous meetings reached consensus that a deep hole through a complete section of fast-spread ocean crust is a renewed priority for the ocean lithosphere community. The scientific rationale for drilling a MoHole in fast-spread crust is developed in the workshop reports (available online), and most thoroughly articulated in the 2007 IODP Mission Moho drilling proposal (IODP Prop 719MP; <http://www.missionmoho.org>; see also Appendix 1).

The 2010 MoHole workshop had two interconnected objectives:

- To initiate a roadmap for the technology development and the project implementation plan that are necessary to achieve the deep drilling objectives of the MoHole project;
- To identify a limited number of potential MoHole sites in the Pacific (i.e., in fast-spread crust), where the scientific community will focus geophysical site survey efforts over the next few years. Selecting drilling sites is essential to clearly identify the range of water depths, drilling target depths and temperatures that we anticipate, and better define the technology required to be developed and implemented to drill, sample and log the MoHole.

The MoHole workshop discussions were held jointly between ocean lithosphere specialists, marine geophysicists, and drilling engineers, with the aim of identifying the best possible sites in fast-spread crust given our current geological, geophysical and technological knowledge and expertise, and to list the items that will need to be considered in a technology roadmap for the MoHole. This report summarizes these discussions, and provides the starting point to initiate the MoHole project.

## 2. Scientific Rationale for the MoHole, a Crustal Journey and Mantle Quest

The Moho is the fundamental boundary within the upper part of our planet, yet we have little knowledge of its geological meaning. New deep drilling technology now make it possible to fulfill scientists' long term aspiration to drill completely through intact oceanic crust, through the seismically defined Moho, and then a significant distance (~500 m) into the upper mantle. Our scientific goals can be divided into the following principal tightly interconnected threads:

- What is the physical nature of the Mohorovičić Discontinuity?, and what is the geological nature of this boundary zone?
- How is the (lower) oceanic crust formed at the mid-ocean ridges, and what processes influence its subsequent evolution? What are the geophysical signatures of these magmatic, tectonic, hydrothermal, and biogeochemical chemical processes?
- What can we infer about the global composition of the oceanic crust, and what are the magnitudes of interactions with the oceans and biology and their influence on global chemical cycles?
- What are the limits of life, and the factors controlling these limits? How do the biological community compositions change with depth, and the evolving physical and chemical environments through the oceanic crust?
- What is the physical and chemical nature of the uppermost mantle, and how does it relate to the overlying magmatic crust?

These objectives (figure 1) are comprehensively articulated in the 2007 Mission Moho proposal (<http://www.missionmoho.org>; see Appendix 1), as well as in the 2006 "Mission Moho" workshop report (<http://www.iodp.org/mission-moho-workshop>; Christie et al., 2006) and the 2009 "Magma, Melting, Fluids and Life" meeting report (<http://www.interridge.org/WG/DeepEarthSampling/workshop2009>; Teagle et al., 2009), and are briefly summarized below.

### 2.1. The Mohorovičić Discontinuity

In the oceans, the Moho is commonly a bright seismic reflector at 5 to 8 km depth and marks an abrupt increase in P-wave seismic velocity ( $V_p$ ) to values in excess of 8 km/s. It is generally assumed that the Moho also represents the boundary between mafic igneous rocks crystallized from magmas that form the crust and residual peridotites of the upper mantle. However, this interpretation has never been tested. Alternatively, there are geologically valid scenarios where the Moho might delineate the boundary between mafic and ultramafic cumulate rocks within the crust, or it may be located below serpentinized peridotites that were previously part of the mantle. Observations and sampling of the Moho, the petrologic crust-mantle boundary and the rocks of the upper mantle are fundamental to understanding the geodynamics and chemical differentiation of our planet. A foremost goal is to reconcile geophysical imaging of the Moho with direct geological observations of cores and downhole measurements; e.g., is the Moho in our study region a sharp compositional boundary or a transition zone of significant thickness (~100s of m)?






Figure 1: Schematic cross-section of fast-spreading crust with anticipated MoHole penetration. The thicknesses of sediment, lavas and sheeted dike complex are taken from ODP/IODP Hole 1256D (Teagle et al., 2006). Top photograph: sheeted dike complex / gabbro contact in Hole 1256D. Predicted physical/chemical profiles in the crust: original figure from Rosalind Coggon (NB : the predicted Mg# profile for a sheeted sill model assumes that on average deeper sills will be more primitive and tend to feed shallower ones, the intrusion of primitive sills at shallow levels cannot be excluded though); lower crust accretion models: after Korenaga and Kelemen (1998). Bottom microphotograph: Mantle peridotite xenolith from French Polynesia (Tommasi et al., 2004). Figure by Benoît Ildefonse.

## 2.2. Formation of the lower crust

On the road to the Moho, we will make paradigm-testing observations of the lower oceanic crust and the deep magmatic, tectonic and hydrothermal processes that occur at the mid-ocean ridges. Our principal target will be intact ocean crust formed at a fast-spreading ridge, which should be relatively laterally uniform, and where we have well-developed theoretical models of crustal accretion that can be tested by drilling. Is the lower oceanic crust formed from the subsidence of a high-level magma chamber, or are there multiple melt bodies at different levels within the oceanic crust (or upper mantle) at fast spreading ridges?

Magnetic stripes document the history of ocean crust formation and are the very basis of plate tectonic theory, yet we have little information on what contribution the lower crust has to this fundamental signature. Similarly, seismic profiling remains the key tool for investigating the deep crust but these regional scale measurements have never been calibrated against core or in situ measurements. It remains challenging to confidently develop geological interpretations from geophysical measurement of the oceanic crust.

### 2.3. Composition and hydration of the ocean crust

A full penetration will provide the first direct estimate of the bulk composition of ocean crust critical for Earth differentiation models. How deeply do seawater-derived hydrothermal fluids penetrate and how efficient is hydrothermal circulation at heat extraction and chemical alteration? Is fluid flow channeled by major faults or more pervasive? (Fig. 2)




Figure 2. A. Schematic architecture of a mid-ocean ridge flank (not to scale), illustrating parameters that may influence the intensity and style of hydrothermal circulation through the ridge flanks, such as faults, seamounts, basement topography, and impermeable sediments, which isolate the crust from the oceans. Arrows indicate heat (red) and fluid (blue) flow. The yellow dashed line is a hypothetical trajectory of the ~120°C isotherm as ocean crust moves away from the ridge crest. B. The calculated global hydrothermal heat flow anomaly decreases to zero, on average, by 65 Ma. C. The effects of parameters such as basement topography and sediment thickness on the intensity and relative cessations of fluid flow, chemical exchange, and microbial activity remain undetermined. D. Evolution of porosity, permeability, and alteration intensity with age. E. Hypothetical change in microbial community structure with; the depth limit of life increases with crustal age. F. Schematic cross-section of fast-spread crust with anticipated MoHole penetration. Figure by Rosalind Coggon; Damon Teagle, and Kentaro Nakamura.

We distinguish two end-member modes of fluid circulation through oceanic crust, i.e., high-temperature circulation at on-/near-axis and low-temperature circulation at off-axis. Near the ridge axis, seawater penetrating into oceanic crust is heated by high-temperature heat source (magma), resulting in vigorous hydrothermal reactions with surrounding rocks and/or phase separation into vapor and brine at relatively shallow level in the oceanic crust, close to the gabbro/sheeted dike interface. Parts of the high-temperature hydrothermal fluids upwell through oceanic crust and finally venting at the seafloor. Some fraction of the high temperature fluids, perhaps segregated brines, may penetrate deeper into the crust, and possibly the mantle, with fluids physical and chemical properties altered by phase separation and host-rock reactions. To date, the physicochemical characteristics, volumes, and depth of penetration of high-temperature fluids are unknown. The MoHole project will provide the first opportunity to access the samples necessary to address these questions.

Equally important will be the characterization of late stage, off-axis low-temperature fluid interactions that will leave an imprint on rocks as the crust ages and cools away from the ridge axis. These lower temperature reactions may be related to, enhanced by, or promote biological activity.

The knowledge of modes of penetration of the hydrothermal fluids, and of the extent of their interactions with the lithosphere, is required to estimate chemical exchanges with the oceans, as well as to assess the volume/variety of materials transferred to the mantle via subduction.

#### **2.4. Limits and controlling factors of life**

Earth is known as a watery planet that is characterized by the presence of liquid water on its surface. Water plays a key role in the cycling of material between lithosphere and hydrosphere (see section 2.3 above). It is well accepted that liquid water is one of the key ingredient contributing to the emergence and development of life on Earth. Elucidating the extent and characteristics of the deep hydrosphere is crucial to understanding not only chemical exchanges between ocean and oceanic crust but also biological activity in the deep subseafloor.

Understanding the limits of life, and the factors controlling these limits, is one of the most fundamental scientific goals for geo- and biosciences, essential for understanding the origin, evolution, distribution, and future of life on Earth as well as celestial bodies. To date, the limits of life even on our own planet remain poorly defined. The MoHole project provides a unique opportunity to address this limit in the oceanic lithosphere that covers ~60% of our planet (Fig. 2E). Numerous factors may control the limits of life, such as temperature, water activity, salinity, pH, energy and carbon sources. Among these, temperature plays a key role, because organism cannot survive beyond as yet poorly known temperature threshold (~110-120 °C?). The ability of seawater to penetrate into the deep crust or mantle and be available for micro-organisms (e.g., minimum pore space), will also have a strong impact on the distribution of living organisms.

#### **2.5. Physical and chemical nature of the upper mantle**

Direct observations of the mantle will document how magmas are focused from a broad melting region to a narrow zone of crustal accretion beneath mid-ocean ridges. Measurements across the Moho will quantify the tectonic coupling between the crust

and mantle. We presently have little knowledge of the composition and physical state of in situ convecting mantle. A few kilograms of fresh residual peridotite from beneath intact oceanic crust would provide a wealth of new information comparable to the treasure trove obtained from the Apollo lunar samples.




Fig. 3 - Schematic summary of the MoHole scientific objectives

### 3. Geophysical Site Surveys

#### 3.1. Scientific Requirements

The criteria for an ideal deep crustal penetration sites have been discussed in previous meetings and workshops (e.g., Christie et al., 2006). Here, we reformulate these criteria based on site survey discussions held during the 2010 MoHole workshop. The selected target would ideally meet all of the following requirements. Satisfying requirements for Points a-e is essential for success. More flexibility is allowed in meeting Points f-h, which are highly desirable but not essential:

- a. Crust formed at fast spreading rate ( $>40$  mm/yr half rate).
- b. Simple tectonic setting with very low-relief seafloor and smooth basement relief; away from fracture zones, propagator pseudo faults, relict overlapping spreading basins, seamounts, or other indicators of late-stage intraplate volcanism. Connection to the host plate active constructive and destructive boundaries would provide important scientific information.
- c. Crustal seismic velocity structure should not be anomalous relative to current understanding of “normal” fast-spread Pacific crust, indicative of layered structure (Fig. 4).
- d. A sharp, strong, single reflection Moho imaged with Multi-Channel Seismic (MCS) techniques (Fig. 4).
- e. A strong wide-angle Moho reflection (PmP), as observed in seismic refraction data, with distinct and clearly identifiable sub-Moho refractions (Pn).
- f. A clear upper mantle seismic anisotropy.
- g. A crust formed at an original latitude greater than  $\pm 15^\circ$ .
- h. A location with relatively high upper crustal seismic velocities indicative of massive volcanic formations to enable the initiation of a deep drill hole.

#### 3.2. Technological Constraints

Several technological constraints limit the range of potential sites. These constraints are:

- Technology for re-circulating drilling mud (riser or alternative) is currently untested at water depths greater than 3000 m. Depths greater than 4000 or 4500 m may exceed the capabilities of a reliable and affordable system.
- Prior scientific ocean drilling experience is mostly limited to temperatures less than 200 °C. Temperatures higher than  $\sim 250^\circ\text{C}$  may limit choices of drill bits and logging tools, may decrease core recovery, and may increase risk of hole failure, or require substantial re-design of drilling equipment. Based on plate cooling models, crust older than  $\sim 15$ -20 Ma should meet this requirement at Moho depths (Fig. 5).
- Thickness of the crustal section above Moho must be at least a few hundred meters less than the maximum penetration/logging/recovery depth of the drilling system, to allow significant penetration in mantle peridotites. For example, if total drilling depth cannot exceed 6 km in 4000 m of water, the targeted crustal section should have a crustal thickness of  $\sim 5.5$  km, somewhat less than average Pacific crust.
- Target area should be in a region with good weather conditions at least 8 months/year, calm seas, and gentle ocean bottom currents.
- Sediment thickness should be greater than 50 m to support possible riser hardware and other seafloor infrastructure (re-entry cones/uppermost casing strings).



- Targeted area should be close ( $\sim 1000$  km) to major port facilities for logistical practicalities.




Fig. 4 - Top left: Example of multichannel seismic reflection image showing a crustal column over a sharp, strong single Moho reflection at the crust-mantle transition zone (image from Nedimovic et al., 2005). Top middle: Seismic structure (one-dimensional  $V_p$  models) of Pacific crust (Grey: Bounds of average "normal" crust older than 29 Ma (after White et al., 1992); Green: ODP site 1256 (Teagle et al., 2006); Red: ODP site 504 (Swift et al., 2008). Top right: Drilled and cored sections in Holes 504B and 1256D (Alt et al., 1993; Wilson et al., 2006), and anticipated lithostratigraphy of MoHole (from Oman ophiolite, e.g., Nicolas and Boudier, 1991). Bottom: Example of OBS wide-angle seismic record section showing crustal refractions ( $P_g$ ), high-amplitude wide-angle Moho reflections/refractions ( $P_mP$ ), and uppermost mantle refractions ( $P_n$ ) (after Canales et al., 2002). These examples illustrate the characteristics of the MCS Moho reflection, wide-angle OBS seismic data, and P-wave velocity structure that will be required to select the MoHole target area.



### 3.3. Potential Sites

Based on the scientific requirements and technological constraints described above, the workshop participants focused the discussions on three areas in the Pacific Basin: Cocos Plate, Off Southern and Baja California, and Hawaii (Tables 1 and 2; Fig. 6). One of the most important issues to take into consideration is the trade off between seafloor depth and temperature at Moho depths (Fig. 5). Most ocean seafloor subsides below 4000 m by ~25 Ma, whereas at Moho depths of 5-7 km temperatures of 200 °C or less are expected for crustal ages of 17-35 Ma.




Figure 5 - Predicted Temperature as a Function of Age and Depth (Half-space thermal model by D. Wilson). Thermal diffusivity is  $6 \times 10^{-7} \text{ m}^2\text{s}^{-1}$ , the initial mantle temperature is 1340°C, the surface temperature is 0°C. At 6 km, cooling below 200°C occurs after 25 Ma.




Fig. 6 - Bathymetric map showing the three selected areas for large-scale MoHole site survey. [A] Cocos plate region, [B] off Southern/Baja California region, [C] off Hawaii region.

Region	Advantages	Disadvantages
Off Southern/Baja California	<ul style="list-style-type: none"> <li>- Large range of water depth</li> <li>- Modest Moho T</li> <li>- higher latitude</li> </ul>	<ul style="list-style-type: none"> <li>- Few data available</li> <li>- Off-ridge volcanism</li> </ul>
Cocos Plate	<ul style="list-style-type: none"> <li>- Shallowest water depth</li> <li>- Well-known tectonics</li> <li>- Sits within a corridor that includes a complete tectonic plate life cycle</li> </ul>	<ul style="list-style-type: none"> <li>- Highest Moho T</li> <li>- Faster than present-day fastest spreading rate</li> <li>- Near equator</li> </ul>
Hawaii	<ul style="list-style-type: none"> <li>- Lowest T</li> <li>- Nearby a large port</li> </ul>	<ul style="list-style-type: none"> <li>- Deepest water</li> <li>- Near large Hotspot</li> <li>- Close to arch volcanism</li> <li>- Near equator</li> <li>- Lowest end of fast-spreading rates</li> </ul>

Table 1 – Regions of interest for preliminary site survey, with principal advantages and disadvantages

Candidate Site	Location	Half spreading rate (mm/yr)	Crustal Age (Ma)	Inferred Moho-T (°C)	Water Depth (m)	Sediment Thickness (m)	Crustal thickness (km)	Total length to the Moho (km)	Original latitude
off S/Baja California A (DeepTow Site)	31-33N, 125-127W	60	30-27	<200	4300-4500	~100	~5.5?	~10?	~30-33
off S/Baja California B	28-29N, 123-125W	50	27-22	~200	4200-4400	~100	?	?	~28
off S/Baja California C	25-26N, 120-122W	60	27-22	~200	3900-4300	~80	?	?	~30
off S/Baja California D	30.5-31N, 121W	45	20-22	~250	2700-4100	~130	?	?	~25
Cocos A (Site 1256)	6.7N, 91.9W	110	15	>250	3646	250	5,5	8.7-9.2	Near equator
Cocos B (Site 844)	8N, 90.5W	100	17	>250	3414	290	5,5?	8.7-9.2?	Near equator
Cocos C	8.7N, 89.5W	100	19	~250	~3400	~300	5,5?	8.7-9.2?	Near equator
Hawaii	22.9-23.7N, 154.9-155.8W	35-40	79-81	~150	4050-4300	~200	~6?	10-10.5?	Near equator
Hawaii	23.5-23.9N, 154.5-154.8W	35-40	78	~150	4300-4500	~200	~6?	10-10.5?	Near equator

Table 2 - Possible candidate sites, with principal characteristics

### 3.3.1. Cocos Plate (aka 1256 area)

This region encompasses a section of the Cocos Plate off Central America (from Guatemala to northern Costa Rica, Fig. 3A) with lithospheric ages between 15 and 25 Ma (Fig. 7). At its western limit on 15 Ma crust, this area includes the ODP Hole 1256D (Wilson et al., 2006; Teagle et al., 2006), a site of on-going IODP deep drilling into intact ocean crust. MCS (Hallenborg et al., 2003; Wilson et al., 2003) and wide-angle OBS data exists for the 15-17 Ma area in the vicinity of Site 1256.




Figure 7 - Map of magnetic anomalies in IODP Site 1256 area (Cocos plate; after Wilson et al., 2003). A-C boxes correspond to the areas detailed in table 2.

The advantages of this region include:

- Super-fast spreading rate (half-spreading rate 110 mm/yr).
- Water depths < 3650 m.
- Crustal thickness: ~5.0 km.
- Nearly planar, sub-horizontal Moho (based on 3D wide-angle OBS data).
- Drilling experience at Site 1256 demonstrates that here the dike section can be drilled successfully.
- This region sits within a corridor that includes a complete tectonic plate life cycle, making it an excellent candidate for understanding ocean crust evolution from a spreading center to subduction. Structure of the crust within this area can be directly related to processes occurring at the modern East Pacific Rise and the Central American subduction zone.
- Areas of older crust (17-24 Ma) probably have drillable Moho temperatures (~200 °C).
- 12-month weather window, 3-m swell rare.

Disadvantages are:

- Elevated Moho temperature (~250-300 °C) at Site 1256.
- Crust formed near the paleo-equator.
- Faster than present-day fastest spreading rate.

- Available MCS data (cruise EW9903) shows poor Moho reflection. Two reasons for this were discussed: (a) weak reflectivity of the Moho is a natural consequence of crust formed at such super-fast rates; or (b) the data may not be of optimal quality (poorly tuned airgun array? source configuration not optimal for low frequencies?)
- 1.0-1.5 knots whole-water-column tidal currents.

### 3.3.2. Off Southern/Baja California

This region encompasses a section of the eastern Pacific Plate off Southern and Baja California, between  $\sim 20$ - $33^\circ\text{N}$ , and  $\sim 130$ - $118^\circ\text{W}$  (Fig. 3B, Fig. 8). Crustal ages are  $\sim 20$ - $35$  Ma. Very little modern geophysical information exists from this region. The best-studied area is in the northernmost part off San Diego, the “Deep Tow” site at  $32^\circ 25'\text{N}$ ,  $125^\circ 45'\text{W}$  (31-32 Ma; Luyendyk, 1970). Historical data there include deep-tow sidescan and bathymetry, 3.5 kHz profiler, magnetics, and single channel seismics. Spreading rate is 55 mm/yr (half rate), sediment thickness is 40-60 m. No modern geophysical data is available to evaluate crustal structure and the presence/characteristics of Moho.




Figure 8 – Map of magnetic anomalies in the region off southern/Baja California. A-D boxes correspond to the areas detailed in table 2. Figure by Doug Wilson.

Advantages of this area are:

- Water depth varies mostly between 4,000-4,500 m, but some areas are shallower than 4,000 m.
- Modest inferred Moho temperature ( $< 250^\circ\text{C}$ ).
- Crustal thickness: 5.5 km estimated 70 km NW of Deep-Tow site using constant layer modeling (Shor, 1968 communication to Luyendyk, 1970).

- Portions of region are close enough for shore logistics base.

Disadvantages are:

- Most of the region has no opportunity to place results within context of ridge processes/evolution since spreading axis is subducted or (micro) plate reorganization have occurred.
- There is so little sub-sediment seismic information available that it is difficult, at this point, to assess either advantages or disadvantages in this region.

### 3.3.3. Hawaii

This area is located north of Oahu in the flexural arch, where water depths are 4000-4300 m (Fig. 9). The crust is ~80 Ma, and was formed at a half spreading rate of 35-40 mm/yr.




Fig. 9 – Magnetic anomaly (A) and bathymetric (B) maps of the region of interest north of Hawaii. Red boxes correspond to the two areas detailed in table 2.

Advantages of this area are:

- Moho temperatures are inferred to be in the 100-150 °C range.
- Close to major port for logistics support.
- 12-month weather window (consistent trade winds but only episodic storms).

Disadvantages are:

- Original latitude near the equator.
- Many seafloor volcanic fields.
- Crustal structure is potentially affected by hotspot volcanism (underplating? crustal intrusions?)
- Spreading rate at the lowest end of what is considered “fast”.



- Cretaceous age make complicate interpretations of hydrothermal alteration due to major differences from modern oceanic chemistry.

### 3.4. Large Scale Surveys: Finding the Right Project Area

The existing geophysical data at the three potential sites are not sufficient to identify a clear MoHole Project target area. Consensus at the workshop was that the priority of the community should be directed toward conducting large-scale seismic surveys in the three regions that will lead to the identification of a Moho target that meets the requirements indicated above (Fig. 4). These surveys should collect spatially coincident MCS data, wide-angle OBS data, multi-beam bathymetry and gravity. As feasible, heat flow and magnetic anomaly data would be useful. The seismic surveys should include the following characteristics:

MCS surveys:

- 2D profiles both parallel and orthogonal to paleo-spreading direction.
- 6 km or longer streamer
- Powerful airgun array well tuned and configured for broadband source rich in low frequencies.

OBS surveys:

- 2D profiles both parallel and orthogonal to paleo-spreading direction with enough aperture to record Pn first arrivals.
- Instrument spacing in the range of 5-10 km.
- Powerful airgun array well tuned and configured for broadband source rich in low frequencies.

JAMSTEC indicated that in 2011 they will dedicate 3 months of science ship time for large-scale surveys. Time required for obtaining environmental permitting for Hawaii precludes conducting surveys in this area in the short term. Discussions were then focused on prioritizing the 1256 region and the Off Southern/Baja California region. Consensus was to prioritize the Off Southern/Baja California region, because so little is known in this area (where depth/age/logistical criteria are viable) that it is urgent to obtain baseline reconnaissance seismic data to assess if the area can possibly meet the scientific requirements. Two factors contributed to the choice of this region as top priority for initial reconnaissance: crustal ages are greater than near Site 1256 so temperatures are expected to be cooler; the existing data suggest that Moho in the 1256 area may not be associated with a simple, continuous, strong reflector.

Survey(s) for the 1256 area remain unplanned and new reconnaissance profiling was considered second priority since EW9903 data provides some information in that region (Wilson et al., 2003; Hallenborg et al., 2003). However it was emphasized that conducting a small survey in this area with state of the art seismic capabilities for comparing with the EW9903 data will be crucial to assess the reason(s) for their low apparent Moho reflectivity (natural structure or data quality?).

The scope and costs of these surveys are too large to be conducted by a single nation or funding agency; *international collaboration is required.*



### 3.5. Small-Scale Surveys: Detailed Imaging of Project Area and Target

After an appropriate drilling target has been found and selected, the community should conduct detailed seismic surveys in the vicinity of the target. The recommended detailed surveys are:

- 2D very-long offset (+8 km) MCS for high-resolution modeling of velocity (i.e., porosity) structure of the upper-mid crust.
- 2D high-resolution, large aperture (+100 km) (crustal and upper mantle) OBS profiles (instrument spacing of ~500 m) for fine-scale structure and physical properties of the drilling target.
- 3D multi-streamer MCS survey for accurate and geometrically correct imaging of intracrustal reflectors (faults, sills, etc.) and Moho.
- 3D crustal and upper mantle scale OBS survey (instrument spacing of ~5 km) to assess crustal structure and thickness variability and upper mantle velocity structure/anisotropy.

While JAMSTEC is taking the lead for conducting large-scale surveys at the Off Southern/Baja California region, other partners (US, Europe) should make an effort to conduct or contribute to other large- and small-scale surveys.

## 4. Technology & Engineering Development

The technology selection and required engineering development will be key components for the success of the MoHole project. It is important to identify potential issues in drilling and coring engineering from the past and on-going ocean drilling expeditions, and to find solutions to overcome the problems encountered. Engineering efforts must remain closely focused on delivering the samples and observations required to achieve the scientific goals of the MoHole endeavor.

Technology selection process and planning for the key engineering developments should be launched as soon as possible in conjunction with site-survey efforts. To do so, establishing a realistic roadmap, which includes project scoping, development and testing elements all controlled by proper project management, is imperative (see “Key for success” section). In the sections to follow, we summarize discussion on each of these items in the MoHole workshop at Kanazawa.

MoHole target sites are all located in Ultra-deep water of approximately 4000m water depth or beyond, and the drilling depth to achieve the MoHole objectives is estimated to extend more than 6000m below seafloor. To drill an ultra deep borehole in such deep water, the provision for continuous mud circulation is a top priority technology requirement. Conventional IODP drilling has utilized riserless drilling which has, given our current experience, drilling depth limit of approximately 2000 to 3000 m.

A promising candidate for drilling to the Moho is riser drilling which provides a conduit for the mud to be returned to the vessel for cleaning, evaluation and recirculation. The drilling vessel *Chikyu* is currently equipped with a deep riser system for maximum rated water depth of 2500m. Significant engineering development is required to prepare *Chikyu* for riser service in water depth near 4000m. In addition to riser drilling several other technologies are being considered to safely and efficiently drill to the targeted depth.

Major areas requiring engineering consideration include :

- mud circulation,
- logging and coring,
- drill bits and drill string,
- drilling mud, and
- casing/cementing.

### 4.1. Mud Circulation Technology

Drilling mud circulation is essential for ultra deep drilling to reach the mantle. We have examined several candidate systems to be equipped on *Chikyu* :

- Riser and BOP (blow out preventer): include CFRP (carbon fiber reinforced plastic) or other light material riser.
- BOP-less: only riser pipes without BOP.
- Surface BOP and SID: slim riser pipe (casing pipe) and Subsea Isolation Device (SID).
- RMR: Riserless mud recovery with mud circulation pump and mud return line (e.g., Myers, 2008).
- Free Standing riser: conventional Riser and BOP on top of huge buoyant& riser.

During the workshop, high-level evaluations were made on each candidate system (table 3). Further investigation should be made including system feasibility, lead-time, safety and cost.

	Riser and BOP ( CFRP riser or other material)	Surface BOP plus SID(ESG)	RMR	RMR with EGS (SID)	Free Standing Riser	BOP-less
Cost	1	3	3	2	2	3
Risk	3	3	2	3	3	0
Lead time	1	3	2	2	1	2
Flexibility	1	3	3	3	2	2
Feasibility	2	3	3	3	2	2
Environmental / Safety	3	3	2	2	3	0
Heat tolerance	2	2	2	2	2	2
Deployment time	1	3	3	2	1	3
Adaptable to <i>Chikyu</i>	3	2	2	2	1	2
Reentry	2	3	3	2	2	2
System reliability	3	3	2	2	2	2
VIV susceptibility	2	2	3	3	2	1
Existing vs development	1	3	3	2	2	3
Seaworthiness	3	2	2	2	3	2
Max. mud weight	3	2	3	3	3	1
Max. flow rate	3	2	2	2	3	2
Casing size options	3	2	2	2	3	3
Max. pressure	3	3	1	1	3	1
Corrosion resistance	2	3	2	2	2	2
Max water depth	1	3	3	3	2	2
Total	43	53	48	45	44	37

Table 3 – Evaluation of mud circulation systems under consideration for MoHole drilling.  
Evaluation ranking : 3 Premium/ 2 Medium/ 1 Poor.

#### 4.2. Logging/ Coring

High formation temperatures (>150°C) are expected at depth in all the candidate sites. Currently available electronic components in logging tools, seal material/bearing, etc., have temperature limits of approximately 150-180°C, and only a small subset of these tools can tolerate short-term exposure to temperatures up to 250°C.

Thus, with current technology, only a few basic measurements such as borehole fluid temperature can be made beyond 250°C, and full suite of measurements needed for the in-situ petrophysical studies will simply not be possible. Therefore, multiple in-situ temperature profiles of ODP Hole 1256D, a riserless hole in which deepening operations are scheduled for 2011 (IODP Expedition 335), are necessary for more accurate

predictions of geothermal gradients to assist planning of further engineering development.

Operational practices such as pumping relatively cooler drilling fluid continuously will be implemented throughout drilling in Hole 1256D. Cooling of the borehole by fluid circulation during wireline logging is also possible.

New, higher temperature, petrophysical logging tools are currently being developed by a number of research groups (e.g., European project HiTi \_High Temperature Instruments, conducted in the frame of the ICDP project IDDP \_Iceland Deep Drilling Project: <http://www.hiti-fp6.eu>), and it is essential that these developments are success in time for Mohole drilling.

#### 4.3. Drill Bit, Drill String, and Core Barrels

Igneous crustal and upper mantle rocks are much harder than sedimentary formations, and are likely to require long drilling times.

Based on past non-riser drilling experience, core recovery rates and the recovery of different rock types can be highly biased, particularly in the upper extrusive sections and dikes, due to variations in fracturing, brecciation, and hydrothermal alteration. On the other hand, core recovery rates are expected to be as high as 80 -100 % in lower crust (gabbros and peridotites), though drilling/coring may suffer from other factors, such as high temperatures, or large in situ deviatoric stresses at depth.

In hard rock formations, drilling/coring bits will be relatively quickly worn out and susceptible to failure, which will significantly reduce the rates of penetration (ROP), and require frequent bit changes. Thus, specifically designed bits for hard and abrasive rocks must be developed with hard bit teeth (e.g., PDC \_polycrystalline diamond compact), and diamond impregnated cutting. Drillbit seals and bearings must also be able to survive long durations in hostile conditions.

The total anticipated depth of operations in the MoHole from sea level is about 10000 to 11000m, which includes water depth of 4000m + drilling depth of 6000 to 7000m. At this great depth, the drill string weight will be close to its maximum tensile strength under its own weight, leaving only a small margin for over-pull. To provide prudent safety margins, ultra-high tensile strength drill pipe must be fabricated (e.g., S160 grade steel that has larger tensile strength than current S140 and S150).

Conventional core barrels may be used if the temperature is not too high, but core barrels developed for geothermal drilling for IDDP may also work at the temperatures anticipated in a MoHole. Developing a wireline coring version of the Iceland Deep Drilling core barrel may be possible.

CDEX (Center for Deep Earth Exploration, JAMSTEC) is currently developing coring bits and drill pipe for ultra-deep drilling.

#### 4.4. Drilling Mud

High formation temperature will negatively impact drilling muds, leading to mud decomposition. The steep temperature gradient between the bottom of the hole and the seafloor may further exacerbate mud stability problems.

For ultra-deep hard-rock drilling, the development of a high density weighed mud for service in high formation temperature is imperative. Currently, the oil and gas industry routinely uses several types of high temperature muds (e.g., oil-based mud, and synthetic based mud \_SBM). However, using such mud types requires leakage-prevention safeguards to avoid adverse environmental impacts.

Geothermal drilling on land has been conducted in very high formation temperatures (>300°C; e.g. Saito et al., 1998; Skinner et al., 2010). Yet, in most cases using water or simple light weight mud with minimum mud contents is sufficient for the completion of the well.

Ultra-deep drilling at sea will require high-density weighed mud to balance high rock pressure and to prevent hole collapse. However, presently available high density weighted muds can be damaged by high temperatures. An alternative might be to develop lighter weight but higher viscosity muds to lift the cuttings as the formation stability in hard igneous rocks should be much higher than the sedimentary rocks encountered in oilfield drilling.

CDEX has initiated discussions with mud engineering companies concerning the development of a muds to withstand high temperatures up to 250°C.

#### 4.5. Casing/Cementing

Casing strings, greater than 6000m, will be extremely heavy and will need to be suspended by drill pipe and lifting rigs on *Chikyu*. Light-weight casing strings may need to be investigated. Detailed well casing designs are essential to safely reach target depths in deep boreholes. The development of a successful well design requires accurate knowledge of pore pressures and fracture gradients. These can be estimated from seismic data, or from offset well data, and must be conducted prior to the commencement of drilling operations in the main hole.

It is common industry practice to drill a pilot well to develop an optimal casing design, and pore pressure gradients can be determined by vertical seismic profiling (VSP) in the pilot hole. High pore pressure may be less of an issue in the igneous rocks to be drilled in the MoHole. It is recognized that trade offs may have to be made regarding the drilling of a pilot hole since it is likely that drilling two deep holes (i.e., a pilot hole and the MoHole) would be prohibitively expensive. However, if drilling is undertaken in discrete stages, drilling can still directly inform the casing program. Possibly a shallow (2000 to 300 m?) pilot hole could be cored to inform the casing plan for the upper section of the deep hole (See also section 5.3.1).

It is essential, for the casing program to successfully enable ultra-deep drilling, that the casing strings are securely bonded to the borehole walls. Secure cementing is also required for the accurate acquisition of data from long-term data monitoring systems that may be placed in the borehole.

This issue of cement and casing must be investigated further with specialist cement engineering companies to develop high temperature, light-weight cements that will ensure excellent bonding between the casing strings and the borehole walls.

#### 4.6. Assessing the Past Experience in Crustal Drilling

It is important to assess the past experience in crustal drilling for optimizing the engineering development and drilling operations. There are several deep holes in the history of the scientific ocean drilling: ODP Holes 504B, 735B, ODP/IODP Hole 1256D, and IODP Hole U1309D. Each hole had different lithology, drilling/coring conditions and strategies, and sea weather conditions. It is noteworthy that 504B and 735B failed due to hole collapse and/or equipment loss (drill bit, pipe).

Assessment of the causes of these events has been partially undertaken by ODP and IODP, but these investigations are not directly available as a self-consistent resource. Past deep drilling information should be compiled and documented for future reference. We suggest that the future scoping group for the Mohole project should include a team to gather the TAMU reviews and IODP-MI Operation Task Force reports of these holes, as well as the driller's log from Transocean and earlier rig-floor operators (SEDCO; Hughes Petroleum for Glomar Challenger).

#### 4.7. Accumulating Experience in Hard-Rock Drilling

In conjunction with the site-survey efforts and the engineering and technology development, it is imperative for ship crews, drilling crews, and scientific technicians on IODP vessels to test capability of the vessels, equipment, and procedures, as well as to accumulate experiences in hard-rock drilling. This will allow the development of drilling/coring strategies, timelines, and protocols for handling the core samples. Prior to the MoHole drilling, thus, it would be prudent to bring *Chikyu* outside of Japan territorial waters to gain valuable drilling experience in ocean crust. The location for this drilling should be based on excellent scientific proposals, some of which have already been developed by the ocean lithosphere drilling community and undergone peer-review (e.g., ocean detachments in slow spreading environment where the lower crust is exposed in relatively shallow water depth). There will be significant return to the MoHole project from the collection of physical property data during ocean crust drilling, allowing to further estimation of borehole stability conditions related to formation types and other factors (e.g., pore pressures, thermal cracking). Such information will provide useful input for the engineering/technology roadmap.



## 5. Operations

Drilling the MoHole will be a challenging enterprise requiring years of detailed preparation, planning, and engineering. The depth of the required borehole is not as deep as other boreholes previously drilled. In fact many boreholes have been drilled to deeper depth including a recently drilled industry hole to over 10 km in the Gulf of Mexico<sup>1</sup>. However, the MoHole is a very different proposition, as it will be cored into hard igneous rocks. To date, the deepest scientific marine borehole is less than 1850 m into igneous basement (ODP Hole 504B; Alt et al., 1993). Operationally, major challenges will be associated with collecting the cored material, making the in-situ measurements, installing casing and keeping the borehole open for successive episodes of deepening in a multiyear, multiphase operation. To gear up for the operations, the following issues must be adequately explored and included in the comprehensive and complete operation plan. At the Kanazawa workshop, it was largely accepted that the drilling to upper mantle in water depths greater than 3650 meters (12000 feet) is feasible. The MoHole initiative is arguably now at the point where the framework for the operations can be constructed since the technology to drill such a hole exists or least has been shown to be feasible.

### 5.1. Professional Planning

This plan must be developed by a dedicated international project office, adequately staffed with scientists, drilling engineers, well designers, engineers, operation specialists, geoscientists, and outreach experts. The current model for managing large projects within the Integrated Ocean Drilling Program is not satisfactory for an undertaking of the scale of the MoHole project. For this project to succeed scientifically, technically, and politically, a central office must be appropriately funded and staffed to take full advantage of all the IODP scientific and technical resources. It is anticipated that all available drilling options, *JOIDES Resolution*, *Chikyu*, as well as Mission Specific Platforms, will be utilized at key stages in the MoHole project to drill boreholes and test newly developed equipment.

### 5.2. Site Characterization Data

Before any detailed plan can be created, the operation group must be supplied with accurate data on the expected borehole temperatures, pressures and lithology to be encountered at all depths of the borehole to be drilled. Fully realizing that there will be some level of uncertainty with this data, such data must be supplied with confidence ranges. Unexpected conditions encountered within the borehole could well terminate deepening operations due to safety concerns, or greatly extended drilling times and costs.

When site characterization is completed together with estimates of expected target depths and sampling plan, detailed design of the well can begin. The key elements of the drilling/coring/downhole measurement are listed below. Each issue must be evaluated to ensure the targets can be reached and to allow an accurate costing for the entire

---

<sup>1</sup> <http://www.bp.com/genericarticle.do?categoryId=2012968&contentId=7055818>  
<http://www.deepwater.com/fw/main/IDeepwater-Horizon-i-Drills-Worlds-Deepest-Oil-and-Gas-Well-419C1.html?LayoutID=6>

project to be determined. This will allow planners to decide which optional elements can be achieved within the available budget.

### 5.3. Well Design Considerations

#### 5.3.1. Pilot Hole Options

The well design of the primary site may require data from a pilot hole or pilot section. The pilot may be a separate hole or it may simply be a pilot section of the main hole. Drilling engineering data from pilot section will be critical in managing the pressure, temperatures and stress within the borehole. Parameters such as mud weights, casing set points will much more accurately be evaluated with the data from the pilot hole. The expected budget constraints may determine whether a separate pilot hole is viable or not.

#### 5.3.2. Well Control and Borehole Pressure Management

The rock types intersected by the borehole drilled to Moho depths are unequivocally expected to be devoid of free gas, overpressured fluid, significant hydrocarbons or other geohazards. Recent events in the Gulf of Mexico may result in future regulatory changes, imposing the use of blow-out prevention (BOP) in mud circulation systems, even in non-hydrocarbon regions. Hence, although a BOP will likely not be needed for well control, the use of a BOP will be considered by the MoHole well planning group.

Mud circulation must be provided to achieve the Mohole objectives and this may be accomplished through the use of a drilling riser or via a riserless mud circulation system. All the requisite support equipment must be considered (see table 3 in section 4.1).

#### 5.3.3. Mud program

The composition of the mud to drill a 6-7 km borehole in crystalline rocks will be vitally important. The engineering decision for which mud circulation option to utilize (see engineering section) will affect the maximum mud weight and viscosity that can be continuously circulated. The mud costs will be a significant portion of the project costs, yet these costs will be minimized since there will be none or only minimal "pump and dump" drilling. Current drilling mud technology is limited to a working temperature of up to 200 to 250°C. Mud will be purchased directly from commercial mud supply companies with representatives onboard during the drilling operation. Personnel will also be required from the mud circulation companies.

#### 5.3.4. Casing design

The casing plan must be created to ensure hole stability to the target depth is achieved with a significant margin of error. The casing plan should be based on widely available casing sizes, and installation hardware. The casing program will hinge on the type of mud circulation system selected and also the type of seafloor safety device (if needed) whether it is conventional blow out preventer (BOP) or a Subsea Isolation Device (SID). If a narrow throat, SID is utilized, then expandable casings must be considered. Regardless of the type of casing, the casing design will need to be extremely comprehensive. The installation of casing the Mohole will be an extremely time consuming and expensive component of the operation. Drill bit selection will be a critical factor for ensuring the successful installation of all the casing strings.

### 5.3.5. Cementing plan

Ensuring the casing is effectively coupled to the formation is another critical element in drilling and coring to the target depth, as well as being essential for the long-term success of borehole sampling/experiments, geophysical logging, and post-drilling observations. Many measurements require complete acoustic coupling for the acquisition of useful data. The cement program must be designed and implemented by cementing professionals with appropriate budget supplied to this element. Personnel from the cementing company will be onboard to oversee the cementing operations.

### 5.3.6. Directional drilling options

Drilling a parallel borehole may be an extremely valuable science tool that, if planned for from the beginning, could be initiated following the completion of the main hole. The design of the direction selection from the main borehole must not impart any risk to the success of the main mission of sampling the lower crust-and mantle rocks. The Integrated Ocean Drilling Program has no experience drilling directional wells, and thus outside consultation and installation expertise must be retained.

## 5.4. Downhole drilling tool selection

The bits and core barrels to be utilized for drilling to the mantle require careful consideration. Bits for drilling versus coring, and the tradeoff between making hole and taking core will require continual discussion and the coring strategy selected may be governed by financial constraints. As drilling in the lower crust approaches the Moho (+500 m), continuous wireline coring will be necessary for the remainder of the well installation. Advances in diamond impregnated coring bits may provide a pathway for improved core recovery. Bottom hole assembly (BHA) hardware choices will also affect hole verticality, thus stabilizers, logging while drilling and deviation avoidance techniques will be critical.

Core barrels and their liners require investigation. With the success of metal liners on MSP expeditions in coral reefs, where the metal liners helped reduce jamming, materials other than plastic should be considered. Most of the drilling in Hole 1256D has been done without plastic liners. Drilling without liners has been common practice in other hard rock sites (Holes 504B, 735B; ODP Leg 209)

## 5.5. Downhole Measurements

### 5.5.1. Drill collar measurements - Logging While Drilling/Measurement While Drilling

Measurements will be made continuously in the drilled borehole, whether it is through logging while drilling (LWD), or coupled with Measurement While Drilling (MWD), where data can be transmitted to the vessel in real-time. In areas of the borehole where continuous wireline coring is required, LWD will not be feasible due to the lack of a provision for a core barrel in current drill collar designs. Logging While Coring (LWC), which allows for simultaneous logging and coring, is possible. Cores can be taken while making resistivity, gamma and resistivity at bit measurements, although such techniques need refinement for hard rock coring (Goldberg et al., 2004). These devices will be very useful in a pilot hole, or just the pilot section of the primary borehole. It must be noted that drill collar measurements afford excellent depth control, but with a trade off of poorer vertical resolution of the tools, which is lower than that of wireline tools.

### 5.5.2. Wireline tools

Following drilling or coring, a full suite of wireline tools will be deployed into the open borehole. Specialty tools including fluid and gas sampling, (bio)geochemical measurements, and even hard rock sidewall coring may be deployed. The temperature of the borehole will be a key limiting factor in the tools that can be deployed. Most commercially available tools have a maximum temperature rating of 175°C, unless new instruments can be developed. Some standard suites of tools can be configured with short-term high temperature exposure ratings up to 250°C. The required temperature tolerances for all required wireline tools need to be explicitly stated, and necessary developments or adaptations compiled to provide necessary lead times. During logging operations, wireline tools and engineers can be called out to the rig as needed, in the same fashion as the oil and gas industry.

### 5.5.3. VSP experiment

Offset vertical seismic profiling will be a critical element of a successful Mohole project. Given the extreme depth of the borehole to be drilled, at least two offset VSP's will be ideally required; one at the estimated half-way point of the well, and one at final depth.

### 5.5.4. Long-term monitoring

Instrumenting the borehole resulting from the Mohole project will eventually become a key, second-stage goal. Thus, the sub-sea equipment and borehole must be constructed to accommodate observatory science. This implies ROV access to the well head and the ability to access the borehole through a BOP or SID.

## 5.6. Logistics

Moving personnel, equipment and supplies routinely, to and from the vessel, will be a critical component of the MoHole operations. Helicopter access and/or workboat access must be provided. Crew changes, provisions, drilling consumables, medical emergencies must all be incorporated in the logistics plans. It was suggested that ideally the drill site should be within 1000 km of major port facilities.

## 6. Keys for success

In this section, we list what are presently considered as the key achievements of a successful MoHole project. These include scientific (essentially sampling strategy) considerations, as well as technology development, industry engagement, and return to the public through outreach and education.

### 6.1. Geophysical surveys and Hole location

The first step to the MoHole Project is the series of geophysical surveys that will be required to identify and characterize an appropriate Project Area within which the drilling will be undertaken. Based on past discussions and consensus (see Christie et al., 2006, Teagle et al., 2009), the MoHole drill site should be in oceanic crust formed at a fast spreading rate, in a region that has aged from the ridge crust with zero perturbation by off-axis volcanism (e.g., seamounts), plume interactions or tectonic disturbance. The Project Area must show a clearly discernible Moho seismic boundary visible in both seismic reflection and refraction records. This drill site will have sufficient sedimentary cover to allow the installation of the (sub-)seafloor infrastructure to enable ultra-deep riser drilling. Relatively high upper crustal seismic velocities, indicating more massive formations in the uppermost volcanic sequences of the crust, will probably be mandatory for successful initiation of a deep drill hole.

### 6.2. Scientific coring, sampling and measurements

Many of our primary scientific goals (see section 2 "scientific rationale") will require continuous core samples.

To be regarded as successful, this experiment must return at least (see also fig. 10):

- Continuous core, including samples of all boundaries, across the region identified by seismic imaging as the Mohorovičić Discontinuity, and the geologic crust-mantle transition from cumulate magmatic rocks to residual peridotites (these may or may not be the same target).
- Continuous coring of the lower 500 m of the mafic and ultramafic cumulate rocks in the oceanic crust.
- Continuous coring of 500 m of peridotites and associated lithologies in the uppermost mantle below the Moho.
- Sufficient cores from intervals of the igneous lower oceanic crust to test models of crustal accretion, melt movement, resolve the geometry and intensity of hydrothermal circulation, and document the limits and activity of the deep microbial biosphere (probably a number of intervals of x100s m of continuous cores).
- Core samples for microbiological studies, using well-established protocols that avoid and/or control chemical and microbiological contaminations from drilling equipments and circulating mud.
- Gas and formation fluid samples from regular intervals within the borehole
- A continuous, comprehensive suite of geophysical logs (wireline, LWD, LWC) and borehole experiments to measure in situ conditions (e.g., temperature and temperature change/gradients) and physical properties, to acquire borehole images, and to identify key geophysical and lithologic regions and transitions (e.g., Layer 2-3 boundary, the Moho) throughout the ocean crust and into the upper mantle.




Figure 10. Technology and financial constraints willing, continuous coring all the way to the Moho and then a significant distance (~500 m) into the uppermost mantle would be the best approach to achieve the scientific goals of this project. However, approaches that mix spot coring (e.g., 10 m coring before bit change every 50 m) with continuous wireline coring may need to be considered. Significant lengths of continuous cores across major lithologic and geophysical transitions will be required and are imperative to answer the fundamental scientific questions posed.

Due to the relatively coarse grain-size of the rocks encountered and the fine scales of expected lithologic/geochemical variation, it is anticipated that lithological records provided by mud/chip logging will be insufficient to address the scientific questions posed. However, a continuous series of mud, cuttings, and gas logs will provide useful



supplementary information in areas of poor or no core recovery, and should be routine throughout the experiment.

In addition to sampling and analyzing rocks, measurements of temperature and chemical compositions of the fluids are required together with biological analyses such as cell counting and DNA/RNA analyses. For example, sidewall core sampling in fracture and breccia zones where crustal fluids pass through may be an effective way to obtain good samples for microbial investigations.

Ultimately, it will be essential to integrate core/log/survey data in a comprehensive synthesis study of the Project Area. Post-drilling studies, such as successive fluid measurements/sampling, *in situ* microbial incubation in the borehole, or VSP experiments should be performed. Therefore, support for post-drilling studies, such as planning CORK installation for long-term borehole fluid measurements and microbial experiments, examination of casing material to avoid metal corrosion that could affect microbial activities, and developing method and tools for measurements and sampling borehole fluids, are important keys to scientific success for the MoHole project.

### 6.3. Broadening the scientific spectrum

The MoHole project is one of the largest and most ambitious scientific endeavors in Earth sciences, comparable to the space exploration missions such as the Apollo Project. This unique challenge will provide precious opportunities to a wide diversity of scientists, engineers and technologists. From a scientific point of view, one of the keys to success of the MoHole project will be the engagement of a broad range of scientific communities. Some, including petrologists, geochemists and geophysicists, have been strongly engaged in science planning for this project so far, but efforts will need to be made to more closely involve other colleagues, such as those from the biological, hydrological or experimental backgrounds. It is important that the MoHole project leaves the door open to new participants, new ideas, and new science.

### 6.4. Drilling technology

At the Kanazawa workshop, it was recognized that several conditions associated with drilling to the MoHole require technological solutions that are yet to be developed. Conditions such as the ultra-deepwater, ultra-deep drilling, the hard dense rock to be drilled, and the extremely high temperature anticipated, are just some of key issues that must be addressed with technology.

Among the many listed items (see section 4, Technology and Engineering Development), the most important one is to develop the technology to drill in the ultra-deepwater close to, or even beyond 4000m. Technologies that are applicable to the MoHole project are now being developed within the oil and gas industry and were presented at the workshop (see also next section). Conversely, some of the technologies required are very specific to scientific drilling, such as logging/coring in high temperature, or drilling the harder crustal and mantle rocks. Therefore, we need to develop such technologies within the IODP community, although these developments may be informed by geothermal projects such as the IDDP in Iceland. CDEX has started to develop some of these technologies using funds outside of the IODP. Key to successfully developing many of the required technologies necessary to complete the MoHole project will be to supplement IODP financial support with other sources of external funding.

The implementation of some of the suggested technologies required to drill the MoHole may require improvements and modifications to the equipment, systems and vessel used for the drilling, most probably the *Chikyu*.

The MoHole undertaking will require more than 10 years to complete, cost hundreds of millions of dollars, and engage a broad international science community. The size and duration of the MoHole effort will require an appropriately funded centralized operations and engineering management group to oversee the project from start to successful completion. This project office will be vital key to success, and in order to succeed, it must be a truly international effort staffed by the best available project managers, engineers, and scientists.

### 6.5. Industrial engagement

Many technologies required to successfully complete the objectives of the MoHole project are "conventional" deep-drilling technologies that are presently being used in the oil and gas industry. At the workshop, technologies for ultra-deep water drilling, such as surface-mounted BOP systems, or the riserless mud recovery system (RMR) were introduced. Such technologies are already being used for deep water drilling in the hydrocarbon industry and are potential candidates for the MoHole project. Other technologies are currently being developed in industry, such as the free-standing riser concept, which would require levels of funding far beyond what is available within the IODP to develop.

Because of the ongoing activities and developments being conducted in ultra-deep water by the oil and gas industry, close and continuous collaboration with industry, and the introduction of new and existing technologies to the MoHole project will be keys to success. For this purpose, it will be necessary to establish a strategy to engage the industry, exchange human resources, and plan joint development work.

Industrial engagement should occur at several levels:

- Ad-hoc collaboration should be continued by inviting oil and gas industry representatives to participate in future planning activities, and to gain their "pro-bono" perspective through the existing IODP science advisory structure and community workshops. This is not a sustainable approach for the safe and complete execution of the MoHole project, but it is useful for scoping technical issues at the beginning of the project.
- Contracting services from planning to execution. This will be the most effective way of benefitting from industry experience in the development and execution of the MoHole drilling initiative.
- Participation of engineers and scientists engaged in the MoHole project to industry workshops, symposiums and technology development forums such as DeepStar (<http://www.deepstar.org>) and RPSEA (<http://www.rpsea.org>). These are important steps in learning about what industry is doing to achieve their technological goals.

### 6.6. Public engagement, outreach, education

In IODP, there have been limited support structures for outreach/education activities mostly from national funding sources, including the Deep Earth Academy and the USIO School of Rocks (Consortium for Ocean Leadership), the ECORD summer schools, and the Core School at the Kochi Core Center (CDEX, Japan). In addition to those initiatives,

there have also been significant outreach activities organized on a case-by-case basis such as the very successful public visits of *Chikyu* in Japan, or the outreach and education team participation on IODP Expedition 327 (Juan de Fuca Hydrogeology 2) in July-August 2010.

One of the key components for the success of the Mohole project will be to improve public support for, and understanding of the scientific goals and excitement of the project. Also we will need to broaden the scientific community engagement by encouraging the involvement of the widest possible range of disciplines (see section 6.5 above). We recommend that engaging the public through outreach and education activities, as well as being pro-active in advertising the project to the wider scientific community and engaging new groups of scientists, be integral parts of the activities carried out by the MoHole project, under the umbrella of IODP and future scientific ocean drilling collaborations. One tool to greatly increase the public and scientific profile of the MoHole project should be a dedicated, multi-language MoHole website, aimed at a range of audiences, that should be implemented as soon as formal scoping activities commence.

## 6.7. Anticipated time table for the MoHole project

### The MoHole timeline

	FY2009	FY2010	FY2011	FY2012	FY2013	FY2014	FY2015	FY2016
General	INVEST Phase II SP writing	1) Kanazawa WS (6/3-5) 2) Sloan WS (9/9-11) <b>Roadmap (Mohole PMT?)</b>	JR Hole 1256D Exp. (April-May) PMT start(?)				<b>Site Decision WS</b> <b>CDEX preparation start(?)</b>	
Characterization of Prospective Site-Data Acquisition	R/V Kilo Moana @Hawaii bathymetry	1) International Site Survey Team	R/V KAIREI off-Mexico	Site 1256 or Hawaii	Site 1256 or Hawaii	Mexico, 1256 or Hawaii 2D 3D seismic(?)	3D seismic(?)	
Data Analysis, & Sample Strategy Development		Sampling Strategies WG		Recon-naissance analysis	data	2D data analysis	3D data analysis	
Technology development		Technology Roadmap & team(?)						

	FY2017	FY2018	FY2019	FY2020	FY2021	FY2022	FY2023
General	CDEX preparation	Start Drilling				Reaching the Mantle	Borehole & Post-drilling research
Science							
Technology & Operation							

## 7. References

- Alt, J.C., Kinoshita, H., Stokking, L.B., et al., 1993. Proc. ODP, Init. Repts., 148: College Station, TX (Ocean Drilling Program). doi:10.2973/odp.proc.ir.148.1993
- Bascom, W.N. 1961. A hole in the bottom of the sea; the story of the Mohole project. Garden City, New York, Doubleday and Company, Inc, 352 p.
- Canales, J. P., G. Ito, R. S. Detrick, and J. M. Sinton, 2002. Crustal thickness along the western Galápagos Spreading Center and the compensation of the Galápagos hotspot swell, *Earth Planet. Sci. Lett.*, 203(1): 311-327. doi:10.1016/S0012-821X(02)00843-9
- Christie, D.M., Ildefonse, B., et al., 2006. Mission Moho - Formation and Evolution of Oceanic Lithosphere. Full workshop Report. Portland, Sept 2006. Available at <http://www.iodp.org/mission-moho-workshop>.
- Goldberg, D., Myers, G., Iturrino, G., Grigar, K., Pettigrew, T., and Mrozewski, S., 2004. Logging-while-coring: first tests of a new technology for scientific drilling. *Petrophysics*, 45(4):328-334.
- Hallenborg, E., A. J. Harding, G. M. Kent, and D. S. Wilson, 2003. Seismic structure of 15 Ma oceanic crust formed at an ultra-fast spreading East Pacific Rise: Evidence for kilometer-scale fracturing from dipping reflectors. *J. Geoph. Res.*, 108(B11), 2532. doi:10.1029/2003JB002400
- Ildefonse, B., Christie, D.M., and Mission Moho Workshop Steering Committee, 2007. Mission Moho workshop : drilling through the oceanic crust to the Mantle. *Scientific Drilling*, 4:11-18. doi:10.2204/iodp.sd.4.02.2007
- Korenaga, J., and Kelemen, P.B., 1998. Melt migration through the oceanic lower crust: a constraint from melt percolation modeling with finite solid diffusion. *Earth and Planetary Science Letters*, 156(1-2): 1-11. doi:10.1016/S0012-821X(98)00004-1
- Luyendyk, B.P., 1970. Origin and History of Abyssal Hills in Northeast Pacific-Ocean. *Geological Society of America bulletin*, 81(8): 2237-2260. doi:10.1130/0016-7606(1970)81[2237:OAHOAH]2.0.CO;2
- Myers, G., 2008. Ultra-deepwater riserless mud circulation with dual gradient drilling. *Scientific Drilling*, 6, 48-51. doi:10.22 04/iodp.sd.6.07.2008
- Nedimovic, M.R., Carbotte, S.M., Harding, A.J., Detrick, R.S., Canales, J.P., Diebold, J.B., Kent, G.M., Tischer, M., and Babcock, J.M., 2005. Frozen magma lenses below the oceanic crust. *Nature*, 436(7054): 1149-1152. doi:10.1038/nature03944
- Nicolas, A., and Boudier, F., 1991. Rooting of the sheeted dike complex in the Oman ophiolite. In: T. Peters, A. Nicolas and R.G. Coleman (Editors), *Ophiolite genesis and evolution of the oceanic lithosphere*. Kluwer Academic Publisher, Dordrecht, pp. 39-54.
- Saito, S., Sakuma, S., and Uchida, T., 1998. Drilling procedures/techniques and test results for a 3.7 km deep, 500°C exploration well, Kakkonda, Japan. *Geothermics*, 16, 462-489. doi:10.1016/S0375-6505(98)00034-0

- Shor, E.N. 1985. A chronology from Mohole to JOIDES. Pp. 391-399 in *Geologists and Ideas; A History of North American Geology*. E.T. Drake and W.M. Jordan, eds. Geol. Soc. Am. Spec. Publ. 4.
- Skinner, A., Bowers, P., Þórhallsson, S., Friðleifsson, G.O., and Guðmundsson, H., 2010. Drilling and operating a core barrel for the Iceland Deep Drilling Project. *GeoDrilling International*, May 2010, 18-21.
- Swift, S., Reichow, M., Tikku, A., Tominaga, M., and Gilbert, L., 2008. Velocity structure of upper ocean crust at Ocean Drilling Program Site 1256. *Geochemistry Geophysics Geosystems*, 9, Q10013. doi:10.1029/2008GC002188
- Teagle, D.A.H., Alt, J.C., Umino, S., Miyashita, S., Banerjee, N.R., Wilson, D.S., and the Expedition 309/312 Scientists, 2006. *Proc. IODP, 309/312: Washington, DC (Integrated Ocean Drilling Program Management International, Inc.)*. doi:10.2204/iodp.proc.309312.2006
- Teagle, D., Ildefonse, B., Blackman, D.K., Edwards, K., Bach, W., Abe, N., Coggon, R., and Dick, H., 2009. *Melting, Magma, Fluids and Life; Challenges for the next generation of scientific ocean drilling into the oceanic lithosphere*. Full Workshop Report. Southampton, July 2009, <http://www.interridge.org/WG/DeepEarthSampling/workshop2009>.
- Tommasi, A., Godard, M., Coromina, G., Dautria, J.M., and Barszczus, H., 2004. Seismic anisotropy and compositionally induced velocity anomalies in the lithosphere above mantle plumes: a petrological and microstructural study of mantle xenoliths from French Polynesia. *Earth Planet. Sci. Lett.*, 227: 539-556. doi:10.1016/j.epsl.2004.09.019
- White, R.S., McKenzie, D., and O'Nions, R.K., 1992. Oceanic crustal thickness from seismic measurements and rare earth element inversions. *J. Geoph. Res.*, 97(B13): 19683-19715. doi:10.1029/92JB01749
- Wilson, D.S., Hallenborg, E., Harding, A.J., and Kent, G.M., 2003. Data report: site survey results from cruise EW9903. In Wilson, D.S., Teagle, D.A.H., Acton, G.D., et al., *Proc. ODP, Init. Repts.*, 206: College Station, TX (Ocean Drilling Program), 1-49. doi:10.2973/odp.proc.ir.206.104.2003
- Wilson, D.S., Teagle, D.A.H., Alt, J.C., Banerjee, N.R., Umino, S., Miyashita, S., Acton, G.D., Anna, R., Barr, S.R., Belghoul, A., Carlut, J., Christie, D.M., Coggon, R.M., Cooper, K.M., Cordier, C., Crispini, L., Durand, S.R., Einaudi, F., Galli, L., Gao, Y.J., Geldmacher, J., Gilbert, L.A., Hayman, N.W., Herrero-Bervera, E., Hirano, N., Holter, S., Ingle, S., Jiang, S.J., Kalberkamp, U., Kerneklian, M., Koepke, J., Laverne, C., Vasquez, H.L.L., MacLennan, J., Morgan, S., Neo, N., Nichols, H.J., Park, S.H., Reichow, M.K., Sakuyama, T., Sano, T., Sandwell, R., Scheibner, B., Smith-Duque, C.E., Swift, S.A., Tartarotti, P., Tikku, A.A., Tominaga, M., Veloso, E.A., Yamasaki, T., Yamazaki, S., and Ziegler, C., 2006. Drilling to gabbro in intact ocean crust. *Science*, 312(5776): 1016-1020. doi:10.1126/science.1126090

## 8. Workshop Agenda

### Thursday, June 3 - Day 1

09:00 - Registration at Kanazawa-Shi Bunka Hall

*09:20 / 10:00 - Workshop Introduction*

- Meeting overview and information (B. Ildefonse, S. Umino)
- Inaugural speech (Shingo Shibata, Director for Deep Sea Research, MEXT)
- Beyond the Moho: scientific justification for a complete penetration of intact in situ ocean crust and into the upper mantle (D. Teagle)

*10:00 / 12:30 - Available site survey data in the Pacific*

- The Trade-off of depth and temperature in choosing a MoHole site (D. Wilson)
- Hawaii site in comparison with IODP site 1256 (N. Seama)
- Site survey data at ODP Site 1256 (D. Wilson)
- "Deep Tow" Site : ~30 Myr Old Crust off Southern California (D. Blackman)
- IFREE seismic studies toward the Mohole project: results and future plans (S. Kodaira)

*13:30 / 14:30 - Current understanding of technological feasibility*

- MoHole Engineering... Seeking solutions inside and outside of IODP (G. Myers)
- *Chikyu* Drilling challenge for the MoHole (S. Kobayashi)
- The *Chikyu* engineering development (M. Yamao)
- Technical Development of Coring System in the Quest to reach the Earth's Mantle (Y. Shinmoto)

*14:30 / 15:00 - Plenary discussion on lectures*

*15:00 / 15:40 - Breakout sessions 1*

- 1 - Ultra deep drilling technology and project planning (including High Temperature/High Pressure drilling and coring) - Chairs : Y. Isozaki, G. Myers and M. Yamao
- 2 - Site selection and site survey plans - Chairs : P. Canales, M. Nedimovic, and D. Blackman

*16:00 / 17:10 - Breakout sessions 2*

- Continuation of discussion on topics 1 and 2

*17:10 / 18:00 - Plenary session*

- Summaries from breakout group chairs & scribes
- Planning, scoping and funding the MoHole within the context of the 10-year renewal cycle (Y. Tatsumi)



## Friday, June 4 - Day 2

Kanazawa-Shi Bunka Hall

09:00 / 10:00 - Drilling and coring at high temperatures

- Iceland Deep Drilling Project (A. Skinner)
- Drilling deep into hot oceanic crust : constraints and challenges (D. Mainprice)

10:00 / 10:50 - Breakout sessions 3

- Continuation of discussion on topics 1 and 2

Breakout group #1 session included presentations from Cameron and Shell representatives :

- Deepwater drilling with Cameron's ESG system & HPHT BOPs (J.E. Kotrla)
- Major Advantages and Disadvantages of the HP Casing Riser with Surface BOP and SID (F.R. Foreman)
- Conventional Drilling Riser Vs. High pressure Drilling Riser (S.F. Shimonek)

11:10 / 12:30 - Breakout sessions 4

Both breakout groups start separately discussions on topic 3

- 3 - Science Measurements: determination of requisite data for project success (e.g., continuous coring Vs. spot coring); in-situ measurements, Core/borehole logging/mud logging/survey integration.

13:30 / 15:40 - Breakout sessions 5

- Continuation of discussions on topic 3

16:00 / 18:00 - Plenary session

- Summaries from breakout groups
- Final discussion; identification of action items

19:00 - Banquet at Kanazawa Excel Hotel Tokyu

## Saturday, June 5 - Day 3

Kakuma Campus, Kanazawa University

10:00 / 13:00 - Writing Committee meeting (Steering Committee + volunteers)

- Drafting of the meeting report
- Listing action items

## 9. Workshop Participants

<b>Natsue Abe</b>	IFREE/JAMSTEC	Japan	abenatsu@jamstec.go.jp
<b>Kozo Akagawa</b>	Cosmos Shoji Co., Ltd.	Japan	akagawa@cosmos-shoji.co.jp
<b>Ryo Anma</b>	University of Tsukuba	Japan	ranma@sakura.cc.tsukuba.ac.jp
<b>Shoji Arai</b>	Kanazawa University	Japan	ultrasa@kenroku.kanazawa-u.ac.jp
<b>Hiroshi Asanuma</b>	Tohoku University	Japan	asanuma@ni2.kankyo.tohoku.ac.jp
<b>Wataru Azuma</b>	CDEX, JAMSTEC	Japan	azumaw@jamstec.go.jp
<b>Neil Banerjee</b>	University of Western Ontario	Canada	neil.banerjee@uwo.ca
<b>Donna Blackman</b>	Scripps Institution of Oceanography	USA	dblackman@ucsd.edu
<b>J. Pablo Canales</b>	Woods Hole Oceanographic Institution	USA	jpcanales@whoi.edu
<b>Georges Ceuleneer</b>	CNRS -Toulouse	France	georges.ceuleneer@ntp.obs-mip.fr
<b>Trevor Falloon</b>	University of Tasmania	Australia	trevor.falloon@utas.edu.au
<b>Fred R. Foreman</b>	In-Spec, Inc.	USA	fred4man@att.net
<b>Gou Fujie</b>	IFREE/JAMSTEC	Japan	fujie@jamstec.go.jp
<b>Patrick Fulton</b>	Oregon State University	USA	pfulton@coas.oregonstate.edu
<b>Nobuo Geshi</b>	Geological Survey of Japan, AIST	Japan	geshi-nob@aist.go.jp
<b>Yumiko Harigane</b>	Kanazawa University	Japan	harigane@staff.kanazawa-u.ac.jp
<b>Nicholas Harmon</b>	University of Southampton	UK	n.harmon@soton.ac.uk
<b>Benoit Ildefonse</b>	CNRS, Universite Montpellier 2	France	benoit.ildefonse@um2.fr
<b>Satoko Ishimaru</b>	Kanazawa University	Japan	jaja@earth.s.kanazawa-u.ac.jp
<b>Osamu Ishizuka</b>	Geological Survey of Japan/AIST	Japan	o-ishizuka@aist.go.jp
<b>Yoshio Isozaki</b>	CDEX/JAMSTEC	Japan	isozaki@jamstec.go.jp
<b>Issa Kagaya</b>	IODP-MI	Japan	ikagaya@iodp.org
<b>Yasuhiro Kato</b>	President of JAMSTEC	Japan	
<b>Yoshihisa Kawamura</b>	IODP-MI	Japan	ykawamura@iodp.org
<b>Mohammed Khedr</b>	Kanazawa University	Japan	khedrzm@earth.s.kanazawa-u.ac.jp
<b>Shomei Kobayashi</b>	CDEX/JAMSTEC	Japan	shomei@jamstec.go.jp
<b>Shuichi Kodaira</b>	IFREE/JAMSTEC	Japan	kodaira@jamstec.go.jp
<b>Johnnie E Kotrla</b>	Cameron International	USA	john.kotrla@c-a-m.com
<b>Yusuke Kubo</b>	CDEX/JAMSTEC	Japan	kuboy@jamstec.go.jp
<b>Hidenori Kumagai</b>	IFREE/JAMSTEC	Japan	kumagai@jamstec.go.jp
<b>Shin'ichi Kuramoto</b>	CDEX/JAMSTEC	Japan	s.kuramoto@jamstec.go.jp
<b>Hans-Christian Larsen</b>	IODP-MI	Japan	hclarsen@iodp-mi-sapporo.org
<b>David Mainprice</b>	CNRS, Université Montpellier 2	France	David.Mainprice@gm.univ-montp2.fr
<b>Charna Meth</b>	Consortium for Ocean Leadership	USA	cmeth@oceanleadership.org

<b>Katsuyoshi Michibayashi</b>	Shizuoka University	Japan	sekmich@ipc.shizuoka.ac.jp
<b>Seiichi Miura</b>	IFREE/JAMSTEC	Japan	miuras@jamstec.go.jp
<b>Eigo Miyasaki</b>	Engineering/JAMSTEC	Japan	miyazakie@jamstec.go.jp
<b>Sumio Miyashita</b>	Niigata University	Japan	miyashit@geo.sc.niigata-u.ac.jp
<b>Gregory Moore</b>	University of Hawaii	USA	gmoore@hawaii.edu
<b>Greg Myers</b>	Consortium for Ocean Leadership	USA	gmyers@oceanleadership.org
<b>Kentaro Nakamura</b>	Precambrian Ecosystem Lab./JAMSTEC	Japan	kentaron@jamstec.go.jp
<b>Mladen Nedimovic</b>	LDEO / Dalhousie University	US/Canada	mladen@dal.ca
<b>Shunji Nishioka</b>	Mitsubishi Heavy Industries, Ltd.	Japan	syunji_nishioka@mhi.co.jp
<b>Jin-Oh Park</b>	University of Tokyo	Japan	jopark@aori.u-tokyo.ac.jp
<b>Betchaida Payot</b>	Kanazawa University	Japan	bdpayot@earth.s.kanazawa-u.ac.jp
<b>Marie Python</b>	Kyoto University	Japan	marie@bep.vgs.kyoto-u.ac.jp
<b>Matthew Schrenk</b>	East Carolina University	USA	schrenkm@ecu.edu
<b>Nobukazu Seama</b>	Kobe University	Japan	seama@kobe-u.ac.jp
<b>Makoto Seki</b>	Japan Petroleum Exploration Co.	Japan	makoto.seki@japex.co.jp
<b>Shingo Shibata</b>	Dr for Deep Sea Research, MEXT	Japan	shibata@mext.go.jp
<b>Steven Shimonek</b>	Cameron International	USA	Steve.Shimonek@c-a-m.com
<b>Yuichi Shinmoto</b>	CDEX/JAMSTEC	Japan	shinmotoy@jamstec.go.jp
<b>Alexander C. Skinner</b>	ACS Coring Services / IDDP	UK	acscs@blueyonder.co.uk
<b>Kiyoshi Suyehiro</b>	IODP-MI	Japan	ksuyehiro@iodp.org
<b>Katsuhiko Suzuki</b>	IFREE/JAMSTEC	Japan	katz@jamstec.go.jp
<b>Ken Takai</b>	SUGAR Project/JAMSTEC	Japan	kent@jamstec.go.jp
<b>Eiichi Takazawa</b>	Niigata University	Japan	takazawa@geo.sc.niigata-u.ac.jp
<b>Akihiro Tamura</b>	Kanazawa University	Japan	kamui@kenroku.kanazawa-u.ac.jp
<b>Yoshihiko Tamura</b>	IFREE/JAMSTEC	Japan	tamuray@jamstec.go.jp
<b>Yoshiyuki Tatsumi</b>	IFREE/JAMSTEC	Japan	tatsumi@jamstec.go.jp
<b>Damon Teagle</b>	NOC, University of Southampton	UK	dat@noc.soton.ac.uk
<b>Masako Tominaga</b>	Woods Hole Oceanographic Institution	USA	mtominaga@whoi.edu
<b>Susumu Umino</b>	Kanazawa University	Japan	sesumin@staff.kanazawa-u.ac.jp
<b>Yoshiyasu Watanabe</b>	Tokai University	Japan	ywata@scc.u-tokai.ac.jp
<b>Douglas Wilson</b>	University of California, Santa Barbara	USA	dwilson@geol.ucsb.edu
<b>Masaoki Yamao</b>	CDEX/JAMSTEC	Japan	yamao@jamstec.go.jp
<b>Mikiya Yamashita</b>	IFREE/JAMSTEC	Japan	mikiya@jamstec.go.jp



# ***The MoHole***

## **A Crustal Journey and Mantle Quest**

### ***Workshop Report***

**Kanazawa, Japan, 3-5 June 2010**



## **Appendices**

## Appendix 1 - Scientific Objectives of Mission Moho proposal (2007)

These pages are excerpted from the Mission Moho proposal that was submitted to IODP in April 2007 (IODP proposal 719MP). Mission Moho was a multi-stage project comprising five successive targets to be drilled using both non-riser and riser technologies. The "MoHole" was planned as the final stage of Mission Moho. Although, no long-term mission has been adopted by IODP, the scientific objectives related to deep drilling in the ocean crust remain essential to our understanding of the Earth. The full proposal is available online (<http://www.missionmoho.org>); here we present the scientific objectives, *as articulated in the mission Moho proposal*, which are fully relevant to the MoHole project.

### *Lead proponents :*

Benoît Ildefonse, CNRS, Montpellier University, France  
 Natsue Abe, JAMSTEC/IFREE, Yokosuka, Japan  
 Peter B. Kelemen, LDEO, Columbia University, USA  
 Hidenori Kumagai, JAMSTEC/IFREE, Yokosuka, Japan  
 Damon A.H. Teagle, NOCS, Southampton, UK  
 Douglas S. Wilson, University of California, Santa Barbara, USA

### *Co-proponents :*

Gary Acton, University of California, Davis, USA  
 Jeffrey C. Alt, University of Michigan, Ann Arbor, USA,  
 Wolfgang Bach, Universität Bremen, Germany  
 Neil R. Banerjee, University of Western Ontario, Canada  
 Mathilde Cannat, CNRS, IPG Paris, France  
 Richard L. Carlson, Texas A&M University, USA  
 David M. Christie, University of Alaska, Fairbanks, USA  
 Rosalind M. Coggon, University of Michigan, Ann Arbor, USA  
 Laurence Coogan, University of Victoria, Canada  
 Robert Detrick, WHOI, Woods Hole, USA  
 Henry J.B. Dick, WHOI, Woods Hole, USA  
 Jeffrey S. Gee, Scripps, UCSD, San Diego, USA  
 Kathryn Gillis, University of Victoria, Canada  
 Alistair Harding, Scripps, UCSD, San Diego, USA  
 Jeffrey A. Karson, Syracuse university, USA  
 Shuichi Kodaira, JAMSTEC/IFREE, Yokosuka, Japan  
 Juergen Koepke, Universität Hannover, Germany  
 John MacLennan, University of Cambridge, UK  
 Jinichiro Maeda, Hokkaido University, Japan  
 Christopher J. MacLeod, Cardiff University, UK  
 D. Jay Miller, IODP, TAMU, College Station, USA  
 Sumio Miyashita, Niigata University, Japan  
 James H. Natland, University of Miami, USA  
 Toshio Nozaka, Okayama University, Japan  
 Mladen Nedimovic, Dalhousie University, Canada  
 Yasuhiko Ohara, Hydrogeology And Oceanography Dept of Japan  
 Kyoko Okino, Tokyo University, Japan  
 Philippe Pezard, CNRS, Montpellier University, France  
 Eiichi Takazawa, Niigata University, Japan  
 Takeshi Tsuji, JAMSTEC/IFREE, Yokosuka, Japan  
 Susumu Umino, Shizuoka University, Japan

## 1. Primary Motivation for Mission Moho

The formation, evolution and recycling of oceanic lithosphere is the dominant process in the chemical differentiation and physical evolution of our planet. This process encompasses the transfer and transformation of material and energy from Earth's mantle to the crust and from the crust, to the ocean and atmosphere. Independent of sunlight, the evolving ocean crust supports life in unique subsurface and seafloor habitats that may resemble the conditions that enabled the origin of life. Upon its formation at seafloor spreading centers, the oceanic lithosphere records geomagnetic field variability, providing the basis for geomagnetic polarity timescales, plate reconstructions, and estimates of plate motions. From its formation until it is subducted back into the mantle, the oceanic lithosphere interacts with seawater, sequesters surface materials (including water and CO<sub>2</sub>) and recycles them back into the mantle.

Sampling a complete section of crust and shallow mantle was the original motivation for scientific ocean drilling. Mission Moho will be the culmination of a decades-old quest by IODP, ODP and DSDP, since Walter Munk and the AMSOC first proposed Project Mohole in 1957 (e.g., Greenberg, 1974; Shor, 1985). The goal was and remains to understand the composition, structure, and evolution of the oceanic lithosphere through deep scientific drilling, as outlined in the 21<sup>st</sup> Century Mohole Initiative of the IODP Initial Science Plan. This goal has been a core component of planning documents since the inception of scientific ocean drilling (e.g., the "Road to the Moho" chapter in Murray et al., 2000).

The Moho (Mohorovičić Discontinuity) is a seismically imaged, primary acoustic interface that represents the transition between the Earth's crust and the underlying mantle. Whereas oceanic crust is formed by a variety of igneous and metamorphic processes, and has a low seismic velocity, the oceanic mantle has a much higher seismic velocity and is largely composed of residual peridotite. Residual peridotites have lost magma to form the crust during partial melting, and although deformed and recrystallized, have remained relatively solid for more than 4 billion years. Thus the mantle is intrinsically different from the igneous crust.

Uncertainty about the Moho stems from questions about the relationship between the seismic boundary, on the one hand, and the geological crust-mantle transition on the other. Are there "ultramafic" igneous rocks, with high seismic velocities, emplaced below the Moho? Are there hydrothermally altered residual peridotites, with low seismic velocities, above the Moho? Is the Moho an intrusive or tectonic boundary? The floor of a magma body? How sharp is the transition, given the limitations of seismic data analysis ( $\pm 50\text{m}$ )?

In addition to the mysteries surrounding the Moho, we also have major gaps in knowledge about the oceanic lower crust and mantle themselves. How does the igneous crust form, and how does it exchange heat and chemical components with seawater? How is melt transport focused from a broad melting region to a narrow zone of crustal accretion beneath mid-ocean ridges? What is the composition and physical state of the convecting mantle? We have no fresh, *in situ* mantle sample. A few kilograms of fresh residual peridotite from beneath intact oceanic crust would provide a wealth of new information comparable to the treasure trove obtained from the Apollo lunar samples.

To date, the elusive frontier at the Moho, and the enormous mantle domain beneath, have been symbolic, unattainable goals. However, with the recent commissioning of IODP's new riser-drilling vessel, D/V *Chikyu*, the aspirations of generations of Earth



scientists of drilling completely through the oceanic crust to the Moho into the upper mantle ~5-6 km below seafloor, have moved into the realm of technical feasibility.

The wider mid-ocean ridge and oceanic lithosphere community has been involved with the establishment of the plan presented herein via the IODP-MI sponsored Mission Moho workshop held in Portland Oregon in September 2006 (Christie et al., 2006; Ildefonse et al., 2007b; full report: <http://www.iodp.org/mission-moho-workshop>). The basic strategy of Mission Moho is derived from the consensus of about one hundred representatives from the international community. Mandate was given to the writing team and co-proponents to take forward the outline plan on behalf of the wider community. The six lead proponents are willing to serve in the stage 1 core Mission Team. This Team should also comprise seismologists (of which several are included in the co-proponent list), deep drilling engineers (especially riser drilling) from USIO, CDEX and industry, logging tool specialists, and borehole management experts (to improve our control of borehole stability issues).

## 2. Road to the Moho

Since the early 70's when the "Penrose" layered model for the ocean crust (Penrose Conference Participants, 1972) was widely accepted, investigations of the oceanic crust by scientific ocean drilling (Fig. 1), marine geological and geophysical techniques, complemented by ophiolite studies, have expanded our understanding of the architecture of the ocean crust (e.g., Teagle et al., 2004; Dick et al., 2006; Ildefonse et al., 2007c).




Fig. 1 - Summaries of existing scientific drill holes into oceanic crustal and mantle rocks. [A] Penetration for all holes penetrating more than 50 meters into basaltic basement, as a function of spreading rate. Hole 1256D has reached the base of the sheeted dike complex at a shallower depth than the bottom of Hole 504B, which is still in sheeted dikes. Black = DSDP holes, blue = ODP holes, red = IODP holes. After Teagle et al. (2006). [B] Depth of penetration for all ODP and IODP holes deeper than 10 meters below seafloor into gabbroic rocks or residual mantle peridotite. Blue = dominantly gabbro, green = dominantly peridotite, white = no recovery. After Blackman et al. (2006).

Away from transform faults, ocean crust formed at fast spreading rates exhibits a relatively uniform seismic stratigraphy (e.g., Canales et al., 2003). At the ridge crests continuous axial low-velocity zones interpreted to be high level, axial melt lenses are imaged, and well defined Moho reflectors are present within a few kilometers of the axis. This suggests that ocean crust formed at fast spreading rates (>80 mm/yr full rate)

is layered and relatively homogeneous. Although only 20% of modern ridges are fast-spreading ( $>80$  mm/yr), more than 50% of the present day seafloor ( $\sim 30\%$  of Earth's surface), and the great majority of crust subducted into the mantle during the past 200 Ma, was produced at fast spreading ridges. Because of the relatively uniform seismic structure and bathymetry of fast-spreading lithosphere, understanding of crust and mantle genesis and evolution at one site can be extrapolated to a significant portion of Earth's surface with some confidence. Importantly, scientists have well developed theoretical models of contrasting styles of magmatic accretion at fast-spreading ridges. Methods have been proposed to test these model using samples recovered from drilled sections of ocean basement together with complementary studies of ophiolites, in particular the Oman ophiolite. Therefore, the highest priority of Mission Moho is to obtain a continuous sample of the entire crust, the Moho and shallow mantle peridotites, in oceanic lithosphere formed at a fast-spreading rate. Scientific and technological progress towards this ultimate goal will require drilling at several additional sites, in crust formed at both fast- and slow-spreading rates.

Ocean crust formed at slow to ultra-slow rates ( $<40$  mm/yr) is highly heterogeneous both along and across axis particularly towards the end of ridge segments where tectonic extension competes with magmatic accretion (e.g., Karson and Elthon, 1987; Dick, 1989; Cannat et al., 1995, 2006; Canales et al., 2000; Kelemen et al., 2004; Ildefonse et al., 2007a). Such is the variety of accretion on slow spreading ridges that fully characterizing the heterogeneity is beyond the scope of this Mission. However, tectonic windows at fast- and slow-spreading ridges provide exposures of deep crustal rocks, serpentinized upper mantle, and possibly fresh upper mantle peridotites that can be sampled in relatively shallow drill holes using existing technologies.

### 3. Scientific Objectives

By drilling an intact section of ocean crust and upper mantle generated at a fast-spreading ridge, we will address first-order questions about the formation of oceanic crust, the nature of the Moho, and the composition of the Earth's convecting mantle. Specific objectives include:

- Determine the geological meaning of the Moho in a variety of tectonic settings,
- Determine the *in situ* composition of the uppermost mantle, its structure and physical properties, and the physics and chemistry of mantle melt migration processes,
- Determine the bulk composition of the oceanic crust to establish the relationship between the lavas that erupt at the seafloor and the melts that separated from their mantle sources,
- Understand the extent and intensity of hydrothermal exchange between the ocean crust and seawater and estimate the chemical flux returned to the mantle by subduction,
- Determine the mode of magmatic, crustal accretion at fast spreading mid-ocean ridges. What are the size and architecture of mid-ocean ridge magma chambers responsible for the construction of the lower ocean crust?,
- Determine the linkages and feedbacks between magma intrusion, hydrothermal circulation and tectonic activity,
- Calibrate regional seismic measurements against core samples and borehole experiments,

- Understand the origin of marine magnetic anomalies and quantify the contribution of lower crustal rocks to the magnetic signature of the ocean crust,
- Establish the limits of life in the ocean lithosphere.

Addressing these objectives requires sampling and logging in deep and so-far unexplored parts of the ocean lithosphere. Specific science questions and working hypotheses to be tested are summarized below. Details of the individual components of Mission Moho will be fully developed in the proposals hosted beneath the Mission Moho umbrella.

### 3.1. What is the geological meaning of the Moho and seismic layers?

Understanding the seismic structure of the ocean lithosphere requires calibration of remotely obtained regional geophysical data against physical properties and petrological measurements of geological samples. There is a well-established terminology for seismic layering in fast-spread oceanic crust. Layer 1 is locally absent, but present where sediment thickness exceeds a few tens of meters, and has  $V_p < 3$  km/s. Layer 2 is a band with a high gradient in  $V_p$  with depth, ranging from  $\sim 3$ -5 to  $\sim 6.7$  km/s, and Layer 3 is a band with nearly uniform  $V_p$  ranging from  $\sim 6.7$  to  $\sim 7.1$  km/s. The Layer 2/3 boundary is an inflection point, between seismic velocities that increase with depth through Layer 2 and nearly uniform seismic velocity through Layer 3. Below Layer 3, in the “classical” Pacific seismic profiles a sharp transition from  $\sim 7$  to  $\sim 8$  km/s occurs within  $< 500$  m. By analogy with ophiolites and geological samples from the Pacific, these layers are commonly interpreted as pelagic sediment (Layer 1), lavas and fractured, sheeted dikes (Layer 2), intact sheeted dikes and plutonic rocks (Layer 3), and residual mantle peridotite  $\pm$  ultramafic plutonic rocks (below Layer 3).

The only site where geological samples have been recovered from intact oceanic crust at the depth of the Layer 2/3 seismic boundary is DSDP/ODP Hole 504B where the inflection in seismic velocity gradient occurs within the sheeted dikes (Detrick et al., 1994; Alt et al., 1996; Carlson, 2001) and appears to be controlled by alteration and/or the nature and density of cracks in the formation, rather than rock type or grain size. Elsewhere, Hole 1256D data suggest that the layer 2/3 seismic transition has not been yet reached, even though the hole extends below the first appearance of gabbros at the base of the sheeted dikes (Teagle et al., 2006). Drilling deeper at Site 1256, and through the Layer 2/3 boundary in the MoHole (if not at Site 1256), will continue to reveal the geological meaning of the seismic layering of the upper ocean crust.

Seismic velocities in the lower oceanic crust are systematically lower than predicted for gabbros (Korenaga et al., 2001; Behn and Kelemen, 2003). This could be related to the presence of cracks and/or alteration phases, and/or to our poor knowledge of lower crustal composition. Direct sampling of Layer 3 will resolve this issue, and restore our ability to interpret Layer 3 velocities in terms of geologically significant rock properties.

The primary goal of Mission Moho is to sample through the base of Layer 3 and the Mohorovičić discontinuity, and into residual peridotites of the upper mantle. In contrast to the classical interpretation of the Moho as the crust-mantle boundary, Hess (1960) posited that the Moho represents a serpentinization front, i.e. a boundary between fresh peridotite and serpentinite. Partially serpentinized peridotites can have densities and velocities identical to (or even lower than) those of fresh gabbros (e.g. Horen et al., 1996; Carlson and Miller, 1997). At fast-spreading ridges, the Moho is generally sharp, which is thought to indicate that the Moho is a lithological contact between gabbro and ultramafic rock. However,  $V_p$  beneath

the oceanic Moho is generally slightly lower than predicted for unaltered peridotite (Shipboard scientific party, 2004). This could indicate ~10% serpentinization, small proportions of gabbroic lenses intruding residual peridotite, or the presence of ultramafic plutonic rocks below the Moho. Until we drill through the Moho beneath fast-spread crust, this question of the petrological significance of the Moho will remain unresolved.

Laboratory-derived velocities of discrete samples cannot reflect large-scale structures within the oceanic lithosphere. Integration of seismic reflection and drilling data will require detailed wireline sonic logging coupled with vertical and multi-ship offset seismic experiments allowing the measurement of regional mantle anisotropy and crustal structure.

At slow-spreading ridges, serpentinized mantle rocks are commonly incorporated into the crust. Drilling through this type of crust, down to fresh peridotites could provide the first fresh, *in situ* mantle samples and test hypotheses regarding the nature of the Moho. Is it: 1) the boundary between the residual upper mantle and the igneous crust, 2) a broader zone of layered ultramafic and mafic rocks, 3) a serpentinization front, or any combination of these three (Fig. 2)? Assessing the role of serpentinization in modifying the seismic signature of the crust and the transition to typical mantle velocities is most conclusively addressed by deep drilling lower ocean crust and upper mantle unroofed at slow rates of oceanic spreading.




Fig. 2 - Various alternative models of crustal architecture and the Moho for the Atlantis Bank (Southwest Indian Ridge), where Hole 735B was drilled (after Dick et al., 2006). [A] After Muller et al. (1997). [B] After Cannat (1996). [C] After Dick et al. (1991).

### 3.2. Obtaining the first fresh samples of the Earth's convecting mantle

Presently there are NO fresh samples of the convecting mantle. Xenoliths, inclusions brought to the surface in lavas, are (a) mainly derived from continental lithosphere, rather than the convecting mantle, and (b) contaminated by interaction with host lavas. This problem is particularly acute for understanding volatile chemical components that are modified by hydrothermal alteration in tectonically exposed samples, or by host lavas in xenoliths. As a result, hypotheses about oxygen fugacity, sulfide composition

and proportion, CO<sub>2</sub>, CH<sub>4</sub>, graphite, H<sub>2</sub>O, Li, B, He and other noble gas characteristics in the “MORB source” –the convecting upper mantle that partially melts to form Mid-Ocean Ridge Basalts– are sustained largely by inference and assertion. Concentrations and isotope characteristics of volatile elements in Earth reservoirs are vital tracers for global chemical cycling. Other components, such as heat-producing elements U and Th, may be concentrated on grain boundaries (Niu, 2004), but we have no fresh grain boundaries to examine.

The nature and length scale of heterogeneity in the mantle source of mid-ocean ridge basalts (MORB) remains controversial and has fundamental implications for the most basic structure of mantle convection: are ocean island basalts fed by hot plumes that cut through the upper mantle MORB source, or is the MORB source replenished from the same mantle material as ocean island basalts? Such problems remain unsolved because mantle source characteristics are largely inferred from lava compositions. Tectonically exposed peridotites sampled by dredging and shallow ocean drilling are plagued by contamination problems due to their highly reactive nature on the seafloor. Drill core from fresh, oceanic upper mantle will place constraints on the extent and scale of Sr, Nd, Pb, Hf and Os isotope vertical variability, independent of the effects of near-surface alteration and deformation.

Similarly, the grain size and deformation history of unaltered oceanic peridotite remains unknown. These parameters, essential for understanding mantle seismic data (e.g., Faul and Jackson, 2005), melt transport in the mantle (e.g., Spiegelman and Kenyon, 1992), or deformation at decreasing temperature caused by corner flow beneath the ridge, can only be addressed through the recovery of fresh samples of *in situ* mantle peridotite.

### 3.3. Melt focusing and extraction beneath mid-ocean ridges

One of the least constrained, fundamental problems in geodynamics is the focusing of mantle melt beneath spreading ridges. Melt is produced in tiny pores along grain boundaries within a region of the upper mantle extending to more than 100 km depth and laterally for 100's of km on either side of the locus of spreading. How is this melt extracted and crystallized to form oceanic crust within a narrow region, a few km wide, as seismically imaged along the East Pacific Rise? Several, well-defined hypotheses have been outlined, but no consensus will emerge without the direct evidence that would be provided by drilling *in situ* upper mantle.

A key element of this problem is to understand and characterize the upwelling path of partially molten mantle peridotite. Is it passive, plate-driven flow (e.g., Langseth et al., 1966; McKenzie, 1967; Bottinga and Allegre 1973, 1976; Sleep, 1975) or active, buoyancy-driven flow (e.g., Rabinowicz et al., 1984; Whitehead et al., 1984; Buck and Su, 1989; Fig. 3)?

For samples obtained more than a few kilometers from the ridge axis, both models predict flow trajectories and lineation approximately perpendicular to the ridge axis, in a nearly horizontal foliation. However, passive vs. buoyancy driven upwelling can be distinguished from the flow kinematics recorded by olivine crystallographic preferred orientations. Active upwelling models predict that outward horizontal flow in the upper mantle is faster than plate velocity, resulting in an inversion of shear senses on a vertical section close to the Moho (Ceuleneer et al., 1988; Nicolas et al., 1988, 1994). This has

been mapped in the Oman ophiolite (Nicolas et al., 1994; Ildefonse et al., 1995) and would be apparent in drill cores.




Fig. 3 - Examples of 2D (perpendicular to ridge axis) numerical models of mantle passive upwelling (left) and active, buoyant upwelling (right). Black curves: melt flow; white curves: solid flow (Spiegelman, 1996).

Another key element is to understand the transport of melt through the mantle peridotite host rocks. Transport may be by diffuse porous flow (e.g., Phipps Morgan, 1987; Spiegelman and McKenzie, 1987), by focused flow in high porosity dissolution channels marked by dunites (e.g., Kelemen et al., 1995a, 1995b, 1997a), by focused flow in high porosity decompaction channels overlain by a permeability barrier at the base of the cold, overlying lithosphere (e.g., Sparks and Parmentier, 1991; Spiegelman, 1993; Ghods and Arkani-Hamed, 2000; Rabinowicz and Ceuleneer, 2005), by focused flow in high porosity shear zones (e.g., Stevenson, 1989; Kelemen & Dick, 1995; Connolly and Podladchikov, 2000; Holtzman et al., 2003), via passive transport within low permeability, partially molten, buoyant diapirs (e.g., Rabinowicz et al., 1984; Whitehead et al., 1984; Buck and Su, 1989), and/or in fractures initiated as a result of overpressure in an interconnected column of buoyant melt overlain by a permeability barrier (e.g., Nicolas 1986, 1990). All of these processes form distinctive geological features, and should be evident in drill core sampling melt transport features in mantle peridotites below the Moho.

Much of what we know about the composition and temperature of the upper mantle, and their global variability, comes from the chemistries of mid-ocean ridge basalts, which are used to infer the composition of the melt that crossed the Moho (e.g., Klein and Langmuir, 1987, McKenzie and Bickle, 1988). However, the majority of MORBs have  $Mg\# \ll 70$  (where  $Mg\# = 100 \times Mg/(Mg + Fe)$  atomic ratio) whereas primitive melts in equilibrium with mantle peridotites should have  $Mg\#$  ranging from  $\sim 70$  to  $78$  (O'Hara, 1968; Langmuir et al., 1982). Thus, we know that melts undergo partial crystallization to produce more primitive cumulates, with  $Mg\# \leq \sim 90$  (equivalent to that of the mantle residues of MORB formation) before the remaining melt is extracted to erupt as MORB. On average, erupted MORBs record about 50% crystallization (see Shipboard Scientific Party, 2004). Primitive cumulate rocks have only rarely been sampled in the oceans, whereas they should be at least as abundant as erupted lavas and sheeted dikes. We



believe that they must comprise much of seismic Layer 3 in fast-spreading crust, and must be abundant somewhere (near ridge segment centers? as intrusions into mantle peridotite?) at slow-spreading ridges, but these hypotheses remain completely untested. In addition, because the nature of the primitive cumulates is unknown, determining the composition of unfractionated melts requires making numerous assumptions. These assumptions can only be tested by having a complete crustal section from which the integrated composition of the entire crust can be determined (e.g., O'Hara, 1982).

### 3.4. Mode(s) of accretion of the lower crust at fast-spreading ridges

Seismic Layer 2A, inferred to be composed mainly of lavas, constitutes less than a sixth of the total crustal thickness in fast-spreading crust, so that the majority of the crust is inferred to be formed from melt that is intruded into the crust from the mantle. The nature of this process is hotly debated. Our understanding has been limited by the difficulties of geophysically imaging and directly sampling the crust, in particular *in situ* gabbroic lower crust. Consequently, the nature of the magma chambers beneath mid-ocean ridges and the magmatic processes that build the lower crust remain virtually unconstrained. Many fundamental questions remain unresolved. For example: how is melt transported from the mantle through the crust? Where do melts fractionate and crystallize? How, and how fast is heat extracted?

Multi-channel seismic (MCS) profiles across active intermediate and fast spreading ridges commonly reveal bright, low velocity reflectors ~1-2 km below the ridge axis, interpreted to be thin (20-100 m thick) axial magma lens (e.g., Morton and Sleep, 1985a; Detrick et al., 1987; Harding et al., 1989; Singh et al., 1998, 2006a; Kent et al., 2000). Melt lenses have also been imaged at or close to Moho depth (Garmany, 1989; Dunn et al., 2001; Crawford and Webb, 2002; Nedimovic et al., 2005; Singh et al., 2006b). These observations, combined with geological and petrological evidence from in-situ ocean crust and the Oman ophiolite, have led to two competing models of lower crustal accretion at fast-spreading mid-ocean ridges:

1. All of the crystallization occurs in a shallow melt lens, and the accumulated crystal residues subside in a "gabbro-glacier" to build the lower crust (e.g., Henstock et al., 1993; Phipps Morgan and Chen, 1993; Quick and Denlinger, 1993; Fig. 4a).
2. Crystallization of lower crustal gabbros occurs partly (Boudier et al., 1996) or essentially *in situ* (Kelemen et al., 1997b; MacLeod and Yaouancq, 2000) via injection of "sheeted sills" (Boudier et al., 1996; Kelemen et al., 1997b; Fig. 4b-c).




Fig. 4 - Schematic drawings of crustal accretion models (after Korenaga and Kelemen, 1998). [a] Gabbro glacier ductile flow model (e.g., Henstock et al., 1993; Quick and Denlinger, 1993). [b] A hybrid model of ductile flow with sill intrusions (e.g., Boudier et al., 1996). [c] "Sheeted sill" model of *in situ* formation of the lower crust by on-axis sill intrusions (e.g., Kelemen et al., 1997b).

These two end-member models have profoundly different implications for the properties of the lower crust, including its composition, the distribution of melt, the extent of deformation, thermal history, and the geometry, temperature and intensity of hydrothermal fluid-rock exchange. Criteria for distinguishing between the two contrasting models are outlined below following tests developed from ophiolite and limited drill core studies. Drill core samples from a Pacific Ocean crustal section are essential to distinguish between these competing models directly, through systematic measurements of compositions, textures, structures, and igneous contacts as a function of depth through the lower crustal gabbro section (Fig. 5).




Figure 5 - Schematic relative variations in the general trends of latent heat release, bulk Mg#, strain rate, cooling rate, hydrothermal fluid flux, fluid temperature, and intensity of high temperature alteration with depth predicted by end-member "gabbro glacier" (with mainly conductive cooling of the lower crust) and "sheeted sill" (with convective cooling of the lower crust) models of crustal accretion.

**Magmatic processes and architecture of the crust:** The igneous stratigraphy and the nature of igneous contacts will be determined, to evaluate whether or not the lower crust comprises individual magma bodies that were intruded into the lower crust, and fractionated and crystallized *in situ*. In the sheeted sill model the bulk crustal composition will become more evolved upwards (Fig 5) and sub-Moho sills (Kelemen et al., 1997b; Korenaga and Kelemen, 1997) are predicted. In contrast, in the gabbro glacier model there will be no change in bulk crustal composition with depth and sub-Moho sills are not expected (Fig 5). In addition to modal layering, vertical chemical variation is observed in ophiolite gabbros (e.g., Pallister and Hopson, 1981; Malpas et al., 1989; Bédard, 1991; Schouten and Kelemen, 2002), and in plutonic rock sections drilled along the mid-ocean ridges (e.g., Dick et al. 1991, 2000; Cannat et al., 1995; Pedersen et al., 1996; Natland and Dick, 1996; Kelemen et al., 2004; Blackman et al., 2006). The nature of this chemical layering can be used to constrain the size of individual crystallization units (Browning, 1984). Magma lenses  $\leq 10$  m thick crystallized layered gabbros in the Troodos ophiolite lower crust (Browning et al., 1989) and in the crust-mantle transition zone of the Oman ophiolite (Korenaga and Kelemen, 1997).

The nature of chemical layering can also be used to place constraints on the mode of melt migration from the mantle through the lower crust, to form shallow gabbros, sheeted dikes, and lavas. Korenaga and Kelemen (1998) showed how reactive porous

flow of melt through chemically layered gabbros would disrupt correlations between mineral compositions formed during crystal fractionation, and gradually smooth vertical chemical variation via diffusion. The scale of measurements undertaken in drill cores is ideal to address such chemical variations.

***Deformation of the ocean crust:*** If the lower crust is built by the subsidence of material from a high level melt lens in a gabbro glacier, increasing strain with depth is predicted (Fig. 5). In contrast, crustal construction by successive sill injections will not produce systematic gradients in strain with depth. Published data sets from the Oman ophiolite are too limited to be conclusive, and show no significant downward trend (Yaouancq and MacLeod, 2000). The intensity of deformation, manifested by crystal shape and lattice preferred orientations, and to some extent by magnetic fabrics (Gee et al., 2004) can be readily assessed in drill core samples.

***Cooling the lower ocean crust:*** As magmas cool and crystallize, both the latent heat of crystallization and specific heat of cooling must be removed. Heat is transferred through the crust by conduction, or the advection of melt, solid material and seawater-derived hydrothermal fluids. Hydrothermal convection removes heat more rapidly than conduction. The distinct distributions of melt intrusion and crystallization with depth implicit in the two end-member models in Figure 4a and 4c yield different distributions of latent heat removal with depth (Fig. 5). Computer simulations that balance the input of magmatic heat from the mantle to the crust with heat sinks provided by conduction, advection, and hydrothermal circulation have been used to test the crustal accretion models, by constraining the input parameters to yield the best fits to geophysical or geological observations (Sleep, 1975; Morton and Sleep, 1985b; Henstock et al., 1993; Phipps Morgan and Chen, 1993; MacLennan et al., 2004). The gabbro glacier model provides the most efficient geometry for hydrothermal heat extraction, as the latent heat and specific heat of cooling can be readily advected from the lid of the melt lens (Henstock et al., 1993), and predicts a decrease in cooling rate with depth, assuming that deep cooling is predominantly by conduction. However, the sheeted sill model (Fig. 4b-c) can also be successfully simulated, provided that vigorous deep hydrothermal circulation occurs near the ridge axis and there is some crystal subsidence from the axial magma lens (MacLennan et al., 2004; MacLennan et al., 2005). The vigor of hydrothermal convection in the lower crust depends on how close to the ridge axis hydrothermal fluids are able to penetrate deeply in the crust, and on permeability. Small changes in permeability may have a huge effect on the resulting thermal structure (Cherkaoui et al., 2003), rendering thermal models uncertain until tested by data. The contrasting distribution, flux and temperatures of hydrothermal fluids predicted by each of the accretion models imply distinct alteration patterns that will be directly observed in drill cores (Fig. 5).

If hydrothermal fluids penetrate sufficiently deep close enough to the ridge axis, they may lower the melting point of newly crystallized gabbro or gabbroic mush and generate more felsic melts (e.g., Koepke et al., 2007), resulting in the intrusion of silicic veins and plutons with distinctive isotope and compositional characteristics. Silicic veins, produced via crystal fractionation or partial melting, are essential to understanding the crustal budget of many geochemically important elements such as U, Th and Pb (e.g., Hart et al., 1999).

The cooling rate of the lower crust, as a function of depth, can be estimated using (i) “geospeedometers” that exploit elemental (Fe, Ca, Li) diffusion rates in olivine, clinopyroxene, and plagioclase (Ozawa, 1986; Coogan et al., 2002, 2005a, 2005b), (ii) analysis of plagioclase crystal size distributions in gabbros (Garrido et al., 2001). Given the potential for small-scale variability in cooling rates, due to fracture-controlled heterogeneous fluid circulation (Coogan et al., 2006), drill core provides the ideal samples for such studies. In addition, the locking in of magnetic polarity in oceanic gabbros at  $\sim 500^{\circ}\text{C}$  can provide strong constraints on cooling rate. Encountering a series of polarity reversals with increasing depth would reveal the record of the blocking isotherm moving deeper in the crust, with time intervals known separately from calibrating the polarity time scale. Uniform polarity over a great depth range would indicate rapid cooling within a time between field reversals.

Well-established petrologic and geochemical techniques can be used to characterize the nature and relative timing of hydrothermal exchange between seawater and the lower crust, the flux of fluid through the crust, and the depth to which fluid penetrates. Mineral geothermometers and cross-cutting vein mineral sequences, coupled with trace element, strontium isotope, and stable isotope compositions of whole rock samples and mineral separates can be used to establish the temperature- and chemical-evolution of the fluids in the lower crust (e.g., Gregory and Taylor, 1981; Manning et al., 1996; Teagle et al., 1998; Bach et al., 2004; Coggon et al., 2004; Gillis et al., 2005), the extent of fluid channeling along fractures and veins (Manning et al., 1996; Banerjee and Gillis, 2001; Nicolas et al., 2003; Bosch et al., 2004; Coogan et al., 2006), and time integrated fluid fluxes (Bickle, 1992; Bickle and Teagle, 1992; Teagle et al., 2003; Gillis et al., 2005).

### **3.5. Crustal aging and chemical fluxes: from mantle to hydrosphere and back again**

The chemical evolution of the oceanic basement does not stop after the crust crystallizes. There is a discernable deficit in conductive heat flow out to 65 Ma on average and some seawater-rock exchange probably occurs in ocean crust of all ages. Geochemical and petrological constraints on water/rock ratios, time integrated fluid, chemical and isotopic fluxes and the nature of hydrothermal alteration of the crust (see section 3.4) will provide essential information on chemical and thermal exchange between the lithosphere and the oceans, key to global geochemical budgets. Stein and Stein (1994) inferred from heat flow data that 33% of the convective cooling of oceanic lithosphere occurs in crust more than ten million years old, so that drilling may actually sample an active, low temperature hydrothermal system. In any case, drilling in crust as old as possible (see section 5.2) will allow better constraints on the role of hydrothermal alteration in controlling the chemical evolution of seawater (e.g., Davis et al., 2003; Nielsen et al., 2006), and the bulk composition of the igneous crust recycled into the mantle in subduction zones.

### **3.6. Probing the limits of life**

The upper oceanic crust is a habitat for microorganisms. Endolithic microbes colonize fractures in glassy basaltic rocks extracting energy and nutrients from the glass by dissolving it, and leaving behind biomarkers that reveal their former presence (e.g., Bach and Edwards, 2003; Banerjee and Muehlenbachs, 2003; Staudigel et al., 2006). Hydrogen and simple organic compounds can be produced abiotically where water interacts with ultramafic rocks in a variety of geotectonic settings, including portions of slow-spreading mid-ocean ridges (e.g.,

McCollom and Seewald, 2001; Kelley et al., 2005). Microbial activity occurring in the sub seafloor biosphere may have a profound impact on processes and chemical fluxes during water-rock reactions but the depth limits of microbiological activity in the oceanic basement have yet to be fully explored.

Exploring and characterizing the sub seafloor biosphere in the ocean crust will be pursued in all oceanic basement holes. Deep drilling will cross chemical and physical boundaries, involving energy, carbon, nutrients and porosity/permeability, which define our current understanding of habitability and may shed light on deep energy sources for microbial communities. An extraordinary diversity of microorganisms exists in high temperature environments ( $>120^{\circ}\text{C}$ ; Kashefi & Lovley, 2003), and this diversity is reflected in enzymes and other molecules. Deep drilling will provide access to rocky habitats at the edge of the life envelope (Bach et al., 2006) and an opportunity to search for novel microorganisms from high-temperature sub seafloor environments, leading to discovery of new compounds with biotechnology, medical and engineering applications. These environments may also be key in the search for primordial microbial communities that may have been the earliest life forms on Earth (e.g., Furnes et al., 2004; Banerjee et al., 2006).

## References

- Alt, J.C., Laverne, C., Vanko, D.A., Tartarotti, P., Teagle, D.A.H., Bach, W., Zuleger, E., Erzinger, J., Honnorez, J., Pezard, P.A., Becker, K., Salisbury, M.H., and Wilkens, R.H., 1996. Hydrothermal alteration of a section of upper oceanic crust in the eastern equatorial Pacific: a synthesis of results from Site 504 (DSDP Legs 69, 70, and 83, and ODP Legs 111, 137, 140, and 148.). In: J.C. Alt, H. Kinoshita, L.B. Stokking and P. Michael (Eds), Proc. ODP, Sci. Results, 148. College Station, TX (Ocean Drilling Program), 417-434.
- Bach, W., and Edwards, K.J., 2003. Iron and sulfide oxidation within the basaltic ocean crust: extent, processes, timing, and implications for chemolithoautotrophic primary biomass production. *Geochimica et Cosmochimica Acta*, 67(20):3871-3887.
- Bach, W., Edwards, K.J., Hayes, J.M., Huber, J.A., Sievert, S.M., and Sogin, M.L., 2006. Energy in the dark: Fuel for life in the deep ocean and beyond. *Eos, Trans., AGU*, 87, 73-78.
- Bach, W., Garrido, C.J., Paulick, H., Harvey, J. and Rosner, M., 2004. Seawater-peridotite interactions: First insights from ODP Leg 209, MAR 15 degrees N. *Geochem. Geophys. Geosys.*, 5: Q09F26 SEP 10 2004.
- Banerjee, N.R., Furnes, H., Muehlenbachs, K., Staudigel, H., and de Wit, M., 2006. Preservation of biosignatures in 3.5 Ga pillow lavas from the Barberton Greenstone Belt, South Africa. *Earth Planet. Sci. Lett.*, 241, 707-722.
- Banerjee, N. R., and K. M. Gillis, 2001, Hydrothermal alteration in a modern suprasubduction zone: The Tonga forearc crust, *J. Geophys. Res.*, 106(B10), 21,737-21,750.
- Banerjee, N. R. and Muehlenbachs, K., 2003, Tuff life: bioalteration in volcanoclastic rocks from the Ontong Java Plateau, *Geochem. Geophys. Geosys.*, 4, 1037, doi:10.1029/2002GC000470.
- Bédard, J.H., 1991. Cumulate Recycling and Crustal Evolution in the Bay of Islands Ophiolite. *J. Geol.*, 99(2): 225-249.
- Behn, M.D. and Kelemen, P.B., 2003. Relationship between seismic P-wave velocity and the composition of anhydrous igneous and meta-igneous rocks. *Geochem. Geophys. Geosyst.*, 4(5), 1041. doi:10.1029/2002GC000393
- Bickle, M.J., 1992. Transport mechanisms by fluid-flow in metamorphic rocks: oxygen and strontium decoupling in the Trois Seigneurs Massif - a consequence of Kinetic dispersion. *Am. Jour. Sci.*, 292: 289-316.
- Bickle, M.J. and Teagle, D.A.H., 1992. Strontium alteration in the Troodos ophiolite: implications for fluid fluxes and geochemical transport in mid-ocean ridge hydrothermal systems. *Earth Planet. Sci. Lett.*, 113: 219-237.
- Blackman, D.K., Ildefonse, B., John, B.E., Ohara, Y., Miller, D.J., MacLeod, C.J., and the Expedition 304/305 Scientists, 2006. Proc. IODP, 304/305: College Station TX (Integrated Ocean Drilling Program Management International, Inc.). doi:10.2204/iodp.proc.304305.2006
- Bosch, D., Jamais, M., Boudier, F., Nicolas, A., Dautria, J.-M. and Agrinier, P., 2004. Deep and high temperature hydrothermal circulation in the Oman ophiolite-Petrological and isotopic evidence. *J. Petrol.*, 45(6): 1181-1208.
- Bottinga, Y. and Allegre, C J, 1973. Thermal aspects of sea-floor spreading and the nature of the oceanic crust. *Tectonophysics*, 18(1-2), 1-17.

- Bottinga, Y. and Allegre, C. J., 1976. Geophysical, petrological and geochemical models of the oceanic lithosphere. *Tectonophysics*, 32(1-2), 9-59.
- Boudier, F., Nicolas, A. and Ildefonse, B., 1996. Magma chambers in the Oman ophiolite : fed from the top or from the bottom ? *Earth Planet. Sci. Lett.*, 144: 239-250.
- Browning, P., 1984. Cryptic variation within the cumulate sequence of the Oman ophiolite: magma chamber depth and petrological implications. In Gass, I.G., Lippard, S.J., and Shelton, A.W. (Eds.), *Ophiolites and Oceanic Lithosphere. Spec. Publ.—Geol. Soc. London*, 13:71–82.
- Browning, P., Roberts, S., and Alabaster, T., 1989. Fine-scale modal layering and cyclic units in ultramafic cumulates from the CY-4 borehole, Troodos ophiolite: evidence for an open system magma chamber, in drillhole CY-4, the Troodos ophiolite, Cyprus. In Gibson, I.L., Malpas, J., Robinson, P.T., and Xenophontos, C. (Eds.), *Cyprus Crustal Study Project: Initial Report, Hole CY-4. Pap.—Geol. Surv. Can.*, 193–220.
- Buck, W.R. and Su, W., 1989. Focused mantle upwelling below mid-ocean ridges due to feedback between viscosity and melting. *Geophys. Res. Lett.*, 16(7): 641-644.
- Canales, J.P., Detrick, R.S., Lin, J., Collins, J.A. and Toomey, D.R., 2000. Crustal and upper mantle seismic structure beneath the rift mountains and across a nontransform offset at the Mid-Atlantic Ridge (35 degrees N). *J. Geophys. Res.*, 105(B2): 2699-2719.
- Canales, J.P., Detrick, R.S., Toomey, D.R. and Wilcock, W.S.D., 2003. Segment-scale variations in the crustal structure of 150-300 kyr old fast spreading oceanic crust (East Pacific Rise, 8 degrees 15 ' N-10 degrees 5 ' N) from wide-angle seismic refraction profiles. *Geophys. J. Int.*, 152(3): 766-794.
- Cannat, M., 1996. How thick is the magmatic crust at slow spreading oceanic ridges? *J. Geophys. Res.*, 101(B2): 2847-2857.
- Cannat, M., Mevel, C., Maia, M., Deplus, C., Durand, C., Gente, P., Agrinier, P., Belarouchi, A., Dubuisson, G., Humler, E. and Reynolds, J., 1995. Thin crust, ultramafic exposures, and rugged faulting patterns at Mid-Atlantic Ridge (22 degrees N). *Geology*, 23: 49-52.
- Cannat, M., Sauter, D., Mendel, V., Ruellan, E., Okino, K., Escartin, J., Combier, V. and Baala, M., 2006. Modes of seafloor generation at a melt-poor ultraslow-spreading ridge. *Geology*, 34: 605-608.
- Carlson, R.L., 2001. The effects of temperature, pressure, and alteration on seismic properties of diabase dike rocks from DSDP/ODP Hole 504B. *Geophys. Res. Lett.*, 28(20): 3979-3982.
- Carlson, R.L. and Miller, D.J., 1997. A new assessment of the abundance of serpentinite in the oceanic crust. *Geophys. Res. Lett.*, 24(4): 457-460.
- Ceuleneer, G., Nicolas, A. and Boudier, F., 1988. Mantle flow patterns at an oceanic spreading centre : the Oman peridotites record. *Tectonophysics*, 151: 1-26.
- Cherkaoui A. S. M., W. S. D. Wilcock, R. A. Dunn, D. R. Toomey, 2003. A numerical model of hydrothermal cooling and crustal accretion at a fast spreading mid-ocean ridge, *Geochem. Geophys. Geosyst.*, 4 (9), 8616, doi:10.1029/2001GC000215
- Christie, D.M., Ildefonse, B., Abe, N., Arai, S., Bach, W., Blackman, D.K., Duncan, R., Hooft, E., Humphris, S.E., and Miller, D.J., 2006. Meeting report. Mission Moho: Formation and Evolution of Oceanic Lithosphere. *Eos, Trans., AGU*, 87 (48), 539.
- Coggon, R.M., Teagle, D.A.H., Cooper, M.J. and Vanko, D.A., 2004. Linking basement carbonate vein compositions to porewater geochemistry across the eastern flank of the Juan de Fuca Ridge, ODP Leg 168. *Earth Planet. Sci. Lett.*, 219(1-2): 111-128.
- Connolly, J.A.D. and Podladchikov, Y.Y., 2000. Temperature-dependent viscoelastic compaction and compartmentalization in sedimentary basins. *Tectonophysics*, 324(3): 137-168.
- Coogan, L.A., Hain, A., Stahl, S. and Chakraborty, S., 2005a. Experimental determination of the diffusion coefficient for calcium in olivine between 900 degrees C and 1500 degrees C. *Geochim. Cosmochim. Acta*, 69: 3683-3694.
- Coogan, L.A., Howard, K.A., Gillis, K.M., Bickle, M.J., Chapman, H., Boyce, A.J., Jenkin, G.R.T. and Wilson, R.N., 2006. Chemical and thermal constraints on focussed fluid flow in the lower oceanic crust. *Am. J. Sci.*, 306(6): 389-427.
- Coogan, L.A., Jenkin, G.R.T. and Wilson, R.N., 2002. Constraining the cooling rate of the lower oceanic crust: a new approach applied to the Oman ophiolite. *Earth Planet. Sci. Lett.*, 199: 127-146.
- Coogan, L.A., Kasemann, S.A. and Chakraborty, S., 2005b. Rates of hydrothermal cooling of new oceanic upper crust derived from lithium-geospeedometry. *Earth Planet. Sci. Lett.*, 240: 415-424.
- Crawford, W.C. and Webb, S.C., 2002. Variations in the distribution of magma in the lower crust and the Moho beneath the East Pacific Rise at 9°-10° N. *Earth Planet. Sci. Lett.*, 203: 117-130.
- Davis, A.C., Bickle, M.J. and Teagle, D.A.H., 2003. Imbalance in the oceanic strontium budget. *Earth Planet. Sci. Lett.*, 211(1-2): 173-187.



- Detrick, R.S., Buhl, P., Vera, E., Mutter, J., Orcutt, J., Madsen, J. and Brocher, T., 1987. Multi-channel seismic imaging of a crustal magma chamber along the East Pacific Rise. *Nature*, 326: 35-41.
- Detrick, R., Collins, J., Stephen, R. and Swift, S., 1994. In situ evidence for the nature of the seismic layer 2/3 boundary in oceanic crust. *Nature*, 370(6487): 288-290.
- Dick, H.J.B., 1989. Abyssal peridotites, very slow spreading ridges and ocean ridge magmatism, in Saunders, A.D., and Norry, M.J., eds., *Magmatism in the Ocean Basins*, Geol. Soc. Spec. Pub. 42, p. 71-105.
- Dick, H.J.B., Meyer, P.S., Bloomer, S., Kirby, S., Stakes, D. and Mawer, C., 1991. Lithostratigraphic evolution of an in-situ section of oceanic layer 3. In: R.P. Von Herzen, P.T. Robinson et al. (Editors), *Proc. ODP, Sci. Results*, 118: College Station, TX (Ocean Drilling Program), 439-540.
- Dick, H.J.B., J.H. Natland, J.C. Alt, W. Bach, D. Bideau, J.S. Gee, S. Haggas, J.G.H. Hertogen, G. Hirsh, P.M. Holm, B. Ildefonse, G.J. Iturrino, B.E. John, D.S. Kelley, E. Kikawa, A. Kingdon, P.J. LeRoux, J. Maeda, P.S. Meyer, J.D. Miller, H.R. Naslund, Y. Niu, P.T. Robinson, J. Snow, R.A. Stephen, P.W. Trimby, H.U. Worm, and A. Yoshinobu. 2000. A long in-situ section of the lower ocean crust: results of ODP Leg 176 drilling at the Southwest Indian Ridge. *Earth Planet. Sci. Lett.* 179:31-51.
- Dick, H.J.B., Natland, J.H., and Ildefonse, 2006. Past and Future Impact of Deep Drilling in the Ocean Crust and Mantle: An Evolving Order Out of New Complexity. *Oceanography*, 19 (4): 72-80.
- Dunn, R.A., Toomey, D.R., Detrick, R.S. and Wilcock, W.S.D., 2001. Continuous mantle melt supply beneath an overlapping spreading centre on the East Pacific Rise. *Science*, 291: 1955-1958.
- Faul, U.H. and Jackson, I., 2005. The seismological signature of temperature and grain size variations in the upper mantle. *Earth Planet. Sci. Lett.*, 234(1-2): 119-134.
- Furnes, H., Banerjee, N.R., Muehlenbachs, K., Staudigel, H., and de Wit, M., 2004, Early life recorded in Archean pillow lavas. *Science*, 304, 578-581.
- Garmany, J., 1989. Accumulations of melt at the base of young oceanic crust. *Nature*, 340: 628-632.
- Garrido, C.J., Kelemen, P.B. and Hirth, G., 2001. Variation of cooling rate with depth in lower crust formed at an oceanic spreading ridge: Plagioclase crystal size distributions in gabbros from the Oman ophiolite. *Geochim. Geophys. Geosyst.*, 2, 10.1029/2000GC000136.
- Gee, J.S., Meurer, W.P., Selkin, P.A. and Cheadle, M.J., 2004. Quantifying three-dimensional silicate fabrics in cumulates using cumulative distribution functions. *J. Petrol.*, 45(10): 1983-2009.
- Ghods, A. and Arkani-Hamed, J., 2000. Melt migration beneath mid-ocean ridges. *Geophys. J. Int.*, 140(3): 687-697.
- Gillis, K.M., Coogan, L.A. and Pedersen, R., 2005. Strontium isotope constraints on fluid flow in the upper oceanic crust at the East Pacific Rise. *Earth Planet. Sci. Lett.*, 232: 83-94.
- Greenberg, D.S. 1971. Mohole : Geopolitical fiasco. Pp. 343-348 In *Understanding the Earth*. I.G. Gass, P.J. Smith, R.C.L. Wilson, and the Open University, eds. Artemis Press.
- Gregory, R.T. and Taylor, H.P.J., 1981. An oxygen isotope profile in a section of Cretaceous oceanic crust, Samail ophiolite, Oman: evidence for  $\delta^{18}\text{O}$  buffering of the oceans by deep (>5km) seawater-hydrothermal circulation at mid-ocean ridges. *J. Geophys. Res.*, 86: 2737-2755.
- Harding, A.J., Orcutt, J.A., Kappus, M.E., Vera, E.E., Mutter, J.C., Buhl, P., Detrick, R.S. and Brocher, T.M., 1989. Structure of young oceanic crust at 13°N on the East Pacific Rise from expanding spread profiles. *J. Geophys. Res.*, 94(B9): 12163-12196.
- Hart, S.R., Blusztajn, J., Dick, H.J.B., Meyer, P.S. and Muehlenbachs, K., 1999. The fingerprint of seawater circulation in a 500-meter section of ocean crust gabbros. *Geochim. Cosmoch. Acta*, 63(23-24): 4059-4080.
- Henstock, T.J., Woods, A.W. and White, R.S., 1993. The accretion of oceanic crust by episodic sill intrusion. *J. Geophys. Res.*, 98(B3): 4143-4161.
- Hess, H.H., 1960. The AMSOC hole to the earth's mantle. *American Scientist*, 48(2), 254-263.
- Holtzman, B. K., N. J. Groebner, M. E. Zimmerman, S. B. Ginsberg, and D. L. Kohlstedt, 2003. Stress-driven melt segregation in partially molten rocks, *Geochim. Geophys. Geosyst.*, 4 (5), 8607, doi:10.1029/2001GC000258
- Horen, H., Zamora, M. and Dubuisson, G., 1996. Seismic waves velocities and anisotropy in serpentinized peridotites from Xigaze ophiolite: Abundance of serpentine in slow spreading ridge. *Geophys. Res. Lett.*, 23(1): 9-12.
- Ildefonse, B., Billiau, S. and Nicolas, A., 1995. A detailed study of mantle flow away from diapirs in the Oman ophiolite. In: R.L.M. Vissers and A. Nicolas (Editors), *mantle and lower crust exposed in oceanic ridges and in ophiolites*. Kluwer, Dordrecht, pp. 163-177.
- Ildefonse, B., Blackman, D.K., John, B.E., Ohara, Y., Miller, D.J., MacLeod, C.J., and the IODP Expeditions 304-305 Scientists, 2007a. Oceanic Core Complexes and Crustal Accretion at Slow-Spreading Ridges. *Geology*, 35, 623-626 ; doi : 10.1130/G23531A.1

- Ildefonse, B., Christie, D.M., and Mission Moho Workshop Steering Committee, 2007b. Mission Moho workshop : drilling through the oceanic crust to the Mantle. *Scientific Drilling*, 4:11-18. doi:10.2204/iodp.sd.4.02.2007
- Ildefonse, B., Rona, P.A., and Blackman, D.K., 2007c. Deep Sampling of the Crust formed at Mid- Ocean Ridges : Scientific Ocean Drilling provides perspective 'in-depth'. *Oceanography*, 20(1) : 22-33.
- Karson, J.A. and Elthon, D., 1987. Evidence for Variations in Magma Production Along Oceanic Spreading Centers - a Critical-Appraisal. *Geology*, 15: 127-131.
- Kashefi K. and Lovley D.R., 2003. Extending the Upper Temperature Limit for Life. *Science*, 301 (5635), 934. doi: 10.1126/science.1086823
- Kelemen, P.B. and Dick, H.J.B., 1995. Focused Melt Flow and Localized Deformation in the Upper-Mantle - Juxtaposition of Replacive Dunite and Ductile Shear Zones in the Josephine Peridotite, Sw Oregon. *J. Geophys. Res.*, 100(B1): 423-438.
- Kelemen, P.B., Hirth, G., Shimizu, N., Spiegelman, M. and Dick, H.J.B., 1997a. A review of melt migration processes in the adiabatically upwelling mantle beneath oceanic spreading ridges. *Philosophical Transactions of the Royal Society of London Series a-Mathematical Physical and Engineering Sciences*, 355(1723): 283-318.
- Kelemen, P.B., Kikawa, E., Miller, D.J., et al., 2004. *Proc. ODP, Init. Repts.*, 209: College Station, TX (Ocean Drilling Program). doi:10.2973/odp.proc.ir.209.2004.
- Kelemen, P., Koga, K. and Shimizu, N., 1997b. Geochemistry of gabbro sills in the crust-mantle transition zone of the Oman ophiolite: implications for the origin of the oceanic lower crust. *Earth Planet. Sci. Lett.*, 146: 475-488.
- Kelemen, P., Shimizu, N. and Salters, V.J.M., 1995a. Extraction of mid-ocean-ridge basalt from the upwelling mantle by focused flow of melt in dunite channels. *Nature*, 375: 747-753.
- Kelemen, P.B., Whitehead, J.A., Aharonov, E. and Jordahl, K.A., 1995b. Experiments on Flow Focusing in Soluble Porous-Media, with Applications to Melt Extraction from the Mantle. *J. Geophys. Res.*, 100(B1): 475-496.
- Kelley, D.S., Karson, J.A., Fruh-Green, G.L., Yoerger, D.R., Shank, T.M., Butterfield, D.A., Hayes, J.M., Schrenk, M.O., Olson, E.J., Proskurowski, G., Jakuba, M., Bradley, A., Larson, B., Ludwig, K., Glickson, D., Buckman, K., Bradley, A.S., Brazelton, W.J., Roe, K., Elend, M.J., Delacour, A., Bernasconi, S.M., Lilley, M.D., Baross, J.A., Summons, R.T. and Sylva, S.P., 2005. A serpentinite-hosted ecosystem: The lost city hydrothermal field. *Science*, 307(5714): 1428-1434.
- Kent, G.M., Singh, S.C., Harding, A.J., Sinha, M.C., Orcutt, J.A., Barton, P.J., White, R.S., Bazin, S., Hobbs, R.W., Tong, C.H. and Pye, J.W., 2000. Evidence from three-dimensional seismic reflectivity images for enhanced melt supply beneath mid-ocean-ridge discontinuities. *Nature*, 406(6796): 614-618.
- Klein, E.M. and Langmuir, C.H., 1987. Global Correlations of Ocean Ridge Basalt Chemistry with Axial Depth and Crustal Thickness. *J. Geophys. Res.*, 92(B8): 8089-8115.
- Koepke, J., Berndt, J., Feig, S.T., Holtz, F., 2007. The formation of SiO<sub>2</sub>-rich melts within the deep oceanic crust by hydrous partial melting of gabbros. *Contrib. Mineral. Petrol.*, 153:67-84. DOI 10.1007/s00410-006-0135-y
- Korenaga, J., Holbrook, W.S., Detrick, R.S. and Kelemen, P.B., 2001. Gravity anomalies and crustal structure at the southeast Greenland margin. *J. Geophys. Res.*, 106(B5): 8853-8870.
- Korenaga, J. and Kelemen, P.B., 1997. Origin of gabbro sills in the Moho transition zone of the Oman ophiolite: Implications for magma transport in the oceanic lower crust. *J. Geophys. Res.*, 102(B12): 27729-27749.
- Korenaga, J. and Kelemen, P.B., 1998. Melt migration through the oceanic lower crust: a constraint from melt percolation modeling with finite solid diffusion. *Earth Planet. Sci. Lett.*, 156: 1-11.
- Langmuir, C.H., Klein, E.M., and Plank, T., 1982, Petrological systematics of mid-ocean ridge basalts: Constraints on melt generation beneath ocean ridges, in Phipps Morgan, J., Blackman, D.K., and Sinton, J.M., eds., *Mantle flow and melt generation*: Washington, DC, AGU, p. 183-280.
- Langseth, M.G. Jr, Le Pichon, X., and Ewing, M., 1966. Crustal structure of the mid-ocean ridges; 5, Heat flow through the Atlantic Ocean floor and convection currents. *J. Geophys. Res.*, 71(22), 5321-5355.
- MacLennan, J., Hulme, T. and Singh, S.C., 2004. Thermal models of oceanic crustal accretion: Linking geophysical, geological and petrological observations. *Geochem. Geophys. Geosyst.*, 5, Q02F25, doi:10.1029/2003GC000605.
- MacLennan, J., Hulme, T. and Singh, S.C., 2005. Cooling of the lower oceanic crust. *Geology*, 33(5): 357-360.
- MacLeod, C.J. and Yaouancq, G., 2000. A fossil melt lens in the Oman ophiolite: implications for magma chamber processes at fast spreading ridges. *Earth Planet. Sci. Lett.*, 176: 357-373.
- Malpas, J., Brace, T., and Dunsworth, S.M., 1989. Structural and petrologic relationships of the CY-4 drill hole of the Cyprus Crustal Study Project, in: I.L. Gibson, J. Malpas, P.T. Robinson, C. Xenophontos (Eds.),

- Cyprus Crustal Study Project: Initial Report, Hole CY-4, Geological Survey of Canada Paper 88-9, pp. 39-67.
- Manning, C.E., Weston, P.E. and Mahon, K.I., 1996. Rapid high-temperature metamorphism of east Pacific rise gabbros from Hess deep. *Earth Planet. Sci. Lett.*, 144(1-2): 123-132.
- McCollom, T.M. and Seewald, J.S., 2001. A reassessment of the potential for reduction of dissolved CO<sub>2</sub> to hydrocarbons during serpentinization of olivine. *Geochimica Cosmochimica Acta*, 65(21): 3769-3778.
- McKenzie, D.P., 1967. The viscosity of the mantle. *Geophys. J. Royal Astr. Soc.*, 14, 297-305.
- McKenzie, D. and Bickle, M.J., 1988. The Volume and Composition of Melt Generated by Extension of the Lithosphere. *J. Petrol.*, 29(3): 625-679.
- Morton, J.L. and Sleep, N.H., 1985a. Seismic reflections from a Lau basin magma chamber. In: D.W. Scholl and T.L. Vallier (Editors), *Geology and offshore resources of pacific island arcs-Tonga region*. Circum-Pacific Council for Energy and Mineral Resources, Earth Science Series, Houston, Texas, pp. 441-453.
- Morton, J.L. and Sleep, N.H., 1985b. A mid-ocean ridge thermal model : constraints on the volume of axial hydrothermal heat flux. *J. Geophys. Res.*, 90(B13): 11345-11353.
- Muller, M.R., Robinson, C.J., Minshull, T.A., White, R.S. and Bickle, M.J., 1997. Thin crust beneath ocean drilling program borehole 735B at the Southwest Indian Ridge? *Earth Planet. Sci. Lett.*, 148(1-2): 93-107.
- Murray, R.W., Schrag, D.P., and Wheat, C.G., 2002. Opportunities in Geochemistry for Post-2003 Ocean Drilling: Tyngsboro, MA (JOI/USSSP).
- Natland, J.H., and Dick, H.J.B., 1996. Melt migration through high-level gabbroic cumulates of the East Pacific Rise at Hess Deep: the origin of magma lenses and the deep crustal structure of fast-spreading ridges. In Mével, C., Gillis, K.M., Allan, J.F., and Meyer, P.S. (Eds.), *Proc. ODP, Sci. Results*, 147: College Station, TX (Ocean Drilling Program), 21-58.
- Nedimovic, M. R., Carbotte, S. M., Harding, A. J., Detrick, R. S., Canales, J. P., Diebold, J. B., Kent, G. M., Tischer, M., and Babcock, J. M., 2005. Frozen magma lenses below the oceanic crust. *Nature*, 436: 1149-1152.
- Nicolas, A., 1986. A melt extraction model based on structural studies in mantle peridotites. *J. Petrol.*, 27: 999-1022.
- Nicolas, A., 1990. Melt extraction from mantle peridotites : hydrofracturing and porous flow, with consequences for oceanic ridge activity. In: M.P. Ryan (Editor), *Magma Transport and storage*. John Wiley & Sons Ltd, pp. 159-173.
- Nicolas, A., Boudier, F. and Ceuleneer, G., 1988. Mantle flow patterns and magma chambers at ocean ridges : evidence from the Oman ophiolite. *Marine Geoph. Res.*, 9: 293-310.
- Nicolas, A., Boudier, F. and Ildefonse, B., 1994. Evidence from the Oman ophiolite for active mantle upwelling beneath a fast-spreading ridge. *Nature*, 370: 51-53.
- Nicolas, A., D. Mainprice, and F. Boudier, 2003. High-temperature seawater circulation throughout crust of oceanic ridges: A model derived from the Oman ophiolites, *J. Geophys. Res.*, 108 (B8), 2371, doi:10.1029/2002JB002094
- Nielsen, S.G., Rehkamper, M., Teagle, D.A.H., Butterfield, D.A., Alt, J.C. and Halliday, A.N., 2006. Hydrothermal fluid fluxes calculated from the isotopic mass balance of thallium in the ocean crust. *Earth Planet. Sci. Lett.*, 251(1-2): 120-133.
- Niu, Y.L., 2004. Bulk-rock major and trace element compositions of abyssal peridotites: Implications for mantle melting, melt extraction and post-melting processes beneath mid-ocean ridges. *J. Petrol.*, 45(12): 2423-2458.
- O'Hara, M.J., 1968. Are ocean floor basalts primary magma? *Nature*, 220(5168), 683-686.
- O'Hara, M.J., 1982. MORB-A Mohole misbegotten ? *Eos, Trans., AGU*, 63 (24), 537.
- Ozawa, K., 1986. Partitioning of elements between constituent minerals in peridotites from the Miyamori ultramafic complex, Kitakami Mountains, Northeast Japan; estimation of P-T condition and igneous composition of minerals. *Journal of the Faculty of Science, University of Tokyo, Section 2: Geology, Mineralogy, Geography, Geophysics*, 21(3), 115-137.
- Pallister, J.S. and Hopson, C.A., 1981. Semail ophiolite plutonic suite ; field relations, phase variation, cryptic variation and layering, and a model of a spreading ridge magma chamber. *J. Geoph. Res.*, 86: 2593-2644.
- Pedersen, R.B., Malpas, J., and Falloon, T., 1996. In Mével, C., Gillis, K.M., Allan, J.F., and Meyer, P.S. (Eds.), *Proc. ODP, Sci. Results*, 147: College Station, TX (Ocean Drilling Program), 3-19.
- Penrose Conference Participants. 1972. Penrose field conference on ophiolites. *Geotimes* 17:24-25.
- Phipps Morgan, J., 1987. Melt Migration beneath Mid-Ocean Spreading Centers. *Geophys. Res. Lett.*, 14(12): 1238-1241.
- Phipps Morgan, J. and Chen, Y.J., 1993. The genesis of oceanic crust: Magma injection, hydrothermal circulation, and crustal flow. *J. Geophys. Res.*, 98: 6283-6297.

- Quick, J.E. and Denlinger, R.P., 1993. Ductile deformation and the origin of layered gabbro in ophiolites. *J. Geophys. Res.*, 98: 14015-14027.
- Rabinowicz, M. and Ceuleneer, G., 2005. The effect of sloped isotherms on melt migration in the shallow mantle: a physical and numerical model based on observations in the Oman ophiolite. *Earth Planet. Sci. Lett.*, 229(3-4): 231-246.
- Rabinowicz, M., Nicolas, A. and Vigneresse, J.L., 1984. A rolling mill effect in asthenosphere beneath oceanic spreading centers. *Earth Planet. Sci. Lett.*, 67: 97-108.
- Schouten, H. and Kelemen, P.B., 2002. Melt viscosity, temperature and transport processes, Troodos ophiolite, Cyprus. *Earth Planet. Sci. Lett.*, 201(2): 337-352.
- Shipboard Scientific Party, 2004. Leg 209 summary. In : Kelemen, P.B., Kikawa, E., Miller, D.J., et al., 2004. *Proc. ODP, Init. Repts.*, 209: College Station, TX (Ocean Drilling Program), doi:10.2973/odp.proc.ir.209.101.2004
- Shor, E.N. 1985. A chronology from Mohole to JOIDES. Pp. 391-399 in *Geologists and Ideas; A History of North American Geology*. E.T. Drake and W.M. Jordan, eds. *Geol. Soc. Am. Spec. Publ.* 4.
- Singh, S.C., Crawford, W.C., Carton, H., Seher, T., Combier, V., Cannat, M., Canales, J.P., Dusunur, D., Escartin, J. and Miranda, J.M., 2006a. Discovery of a magma chamber and faults beneath a Mid-Atlantic Ridge hydrothermal field. *Nature*, 442: 1029-1032.
- Singh, S. C., A. J. Harding, G. M. Kent, M. C. Sinha, V. Combier, S. Bazin, C. H. Tong, P. J. Barton, R. W. Hobbs, R. S. White and J. A. Orcutt, 2006b. Seismic reflection images of the Moho underlying melt sills at the East Pacific Rise, *Nature*, 442, 287-290.
- Singh, S.C., Kent, G.M., Collier, J.S., Harding, A.J. and Orcutt, J.A., 1998. Melt to mush variations in crustal magma properties along the ridge crest at the southern East Pacific Rise. *Nature*, 394(6696): 874-878.
- Sleep, N H, 1975. Formation of oceanic crust; some thermal constraints. *J. Geophys. Res.*, 80(29), 4037-4042.
- Sparks, D.W. and Parmentier, E.M., 1991. Melt Extraction from the Mantle beneath Spreading Centers. *Earth Planet. Sci. Lett.*, 105(4): 368-377.
- Spiegelman, M., 1993. Physics of Melt Extraction - Theory, Implications and Applications. *Ph. Trans. R. Soc. London, Series a-Mathematical Physical and Engineering Sciences*, 342(1663): 23-41.
- Spiegelman, M., 1996. Geochemical consequences of melt transport in 2-D: The sensitivity of trace elements to mantle. *Earth Planet. Sci. Lett.*, 139(1-2): 115-132.
- Spiegelman, M. and Kenyon, P., 1992. The Requirements for Chemical Disequilibrium During Magma Migration. *Earth Planet. Sci. Lett.*, 109(3-4): 611-620.
- Spiegelman, M. and McKenzie, D., 1987. Simple 2-D Models for Melt Extraction at Mid-ocean Ridges and Island Arcs. *Earth Planet. Sci. Lett.*, 83(1-4): 137-152.
- Staudigel, H., Furnes, H., Banerjee, N.R., Dilek, Y. & Muehlenbachs, K., 2006, Microbes and volcanoes: A tale from the oceans, ophiolites and greenstone belts. *GSA Today*, v. 16, No. 10, p. 4-10, doi: 10.1130/GSAT01610A.1.
- Stein, C. A., Stein, S., 1994. Constraints on hydrothermal heat flux through the oceanic lithosphere from global heat flow. *J. Geophys. Res.*, 99, 3081-3096, 10.1029/93JB02222.
- Stevenson, D.J., 1989. Spontaneous Small-Scale Melt Segregation in Partial Melts Undergoing Deformation. *Geophys. Res. Lett.*, 16(9): 1067-1070.
- Teagle, D.A.H., Alt, J.C. and Halliday, A.N., 1998. Tracing the evolution of hydrothermal fluids in the upper oceanic crust: Sr isotopic constraints from DSDP/ODP Holes 504B and 896A. In: R.A. Mills and K. Harrison (Editors), *Modern Ocean-Floor Processes and the Geological Record*. *Geol. Soc. (Lond.) Spec. Pub. Geol. Soc. (Lond.)*, London, pp. 81-97.
- Teagle, D.A.H., Alt, J.C., Umino, S., Miyashita, S., Banerjee, N.R., Wilson, D.S., and the Expedition 309/312 Scientists, 2006. *Proc. IODP, 309/312*: Washington, DC (Integrated Ocean Drilling Program Management International, Inc.). doi:10.2204/iodp.proc.309312.2006
- Teagle, D.A.H., Bickle, M.J. and Alt, J.C., 2003. Recharge flux to ocean-ridge black smoker systems: a geochemical estimate from ODP Hole 504B. *Earth Planet. Sci. Lett.*, 210: 81-89.
- Teagle, D.A.H., Wilson, D.S., Acton, G.D. and ODP Leg 206 Shipboard Science Party, 2004. The "Road to the MoHole" four decades on: Deep drilling at Site 1256. *EOS, Trans. Am. Geophys. Union*, 85: 521,530-531.
- Whitehead, J.A Jr, Dick, H.J.B., Schouten, H.A., 1984. Mechanism for magmatic accretion under spreading centres. *Nature*, 312(5990), 146-148.
- Yaouancq, G. and MacLeod, C.J., 2000. Petrofabric Investigation of Gabbros from the Oman Ophiolite: Comparison between AMS and rock fabric. *Mar. Geoph. Res.*, 21(3-4): 289-305.

## Appendix 2 - Langseth Workshop White Paper (2010)

This white paper was submitted to the NSF "Langseth" workshop (*Challenges and Opportunities in Academic Marine Seismology*, March 22-24, 2010, Incline Village, NV, USA; [http://www.unols.org/meetings/2010/201003mls/201003LW\\_details.html](http://www.unols.org/meetings/2010/201003mls/201003LW_details.html))

### Setting up the Stage for Project MoHole:

#### Seismic Studies of Fast-Spread Ocean Lithosphere

J.P. Canales<sup>1</sup>, M. Nedimović<sup>2</sup>, B. Ildefonse<sup>3</sup>, S. Kodaira<sup>4</sup>, N. Abe<sup>4</sup>, H.-C. Larsen<sup>5</sup>, D. Wilson<sup>6</sup>, D. Blackman<sup>7</sup>, Y. Isozaki<sup>8</sup>, G. Myers<sup>9</sup>, D. Teagle<sup>10</sup>, S. Umino<sup>11</sup>

<sup>1</sup> Woods Hole Oceanographic Institution, USA

<sup>2</sup> Dalhousie University, Canada

<sup>3</sup> Geosciences Montpellier, CNRS, France

<sup>4</sup> Institute for Research on Earth Evolution, JAMSTEC, Japan

<sup>5</sup> IODP Management International, Japan

<sup>6</sup> University of California Santa Barbara, USA

<sup>7</sup> Scripps Institution of Oceanography, USA

<sup>8</sup> Center for Deep Earth Exploration, JAMSTEC, Japan

<sup>9</sup> Consortium for Ocean Leadership, USA

<sup>10</sup> National Oceanography Centre Southampton, UK

<sup>11</sup> Kanazawa University, Japan

The mid-ocean ridges and the new oceanic lithosphere that they create are the principal pathway for energy and mass exchange and physical/chemical interactions between the earth's interior, the hydrosphere, and the biosphere. Bio-geochemical reactions between the oceans and oceanic crust occur throughout its lifetime, and hence the ocean lithosphere records the inventory of global thermal, chemical and biological exchanges. The MoHole, an initiative to drill an ultra-deep hole in an intact portion of oceanic lithosphere, through the crust to the Mohorovičić discontinuity (Moho), and into the uppermost mantle is a long-standing goal of scientific ocean drilling. It remains critical to answer many fundamental questions about the dynamics of the Earth and global elemental cycles; its fundamental goals include:

- Determine the bulk composition of the oceanic crust to establish the chemical links between erupted lavas and primary mantle melts, understand the extent and intensity of seawater hydrothermal exchange with the lithosphere, and estimate the chemical fluxes returned to the mantle by subduction,
- Test competing hypotheses of the ocean crust accretion at fast spreading mid-ocean ridges, and quantify the linkages and feedbacks between magma intrusion, hydrothermal circulation and tectonic activity,
- Determine the geological meaning of the Moho in different oceanic settings,
- Determine the in situ composition, structure and physical properties of the uppermost mantle (and its variability), and understand mantle melt migration,
- Calibrate regional seismic measurements against recovered cores and borehole measurements, and understand the origin of marine magnetic anomalies,
- Establish the depth extent of deep biosphere and hydrological/geobiological processes in the lithosphere.

More detailed scientific rationale for the MoHole can be found in recent workshop reports:

- <http://www.iodp.org/mission-moho-workshop/>
- <http://www.interridge.org/WG/DeepEarthSampling/workshop2009>

A forthcoming workshop to be held in Kanazawa, Japan, in June 2010 (<http://earth.s.kanazawa-u.ac.jp/~Mohole/>) will identify 2-3 potential MoHole sites in the Pacific where the scientific community will focus geophysical site survey and post-drilling research efforts over the next few years. The type, resolving power, and coverage of geophysical data needed for site selection and to accomplish the post-drilling scientific goals will be amply discussed during the workshop.

In particular, seismic surveys are expected to be a fundamental component of MoHole critical to accomplish the goals of this ambitious project. These surveys will be more efficiently conducted through international collaborations involving several platforms of different characteristics. Among these platforms, the *R/V Langseth* will undoubtedly be a key resource because of her state-of-the-art seismic capabilities. We envision *Langseth* contributing to acquisition of one or more of several types of seismic data needed to support Project MoHole, which include:

(1) 3D multichannel seismic (MCS) data. MCS reflection imaging of the oceanic Moho is often degraded by in- and out-of-plane energy scattered by the rough igneous basement [e.g., Kent *et al.*, 1996]. Among the many benefits of 3D MCS data and 3D processing techniques is the possibility of accurately collapsing the scattered wavefield to its source location; therefore significantly improving image quality. In addition, obtaining geometrically accurate images of steeply dipping faults that may cut an entire crustal section (and therefore perhaps affecting the physical properties of the Moho) [e.g., Nedimović *et al.*, 2009] will also require 3D MCS data/processing. The potential benefits of combining 3D MCS with 3D borehole VSP for a project of these characteristics need to be explored.

(2) 2D long-offset (<sup>3</sup>8-km streamer) MCS data. In recent years 2D MCS data collected with 6-km-long hydrophone streamers have resulted in seismic images of the lower oceanic crust, Moho, and sub-Moho structure of unprecedented quality and detail, contributing to a better understanding of the geological processes that form the lower crust and Moho [Canales *et al.*, 2009; Nedimović *et al.*, 2005]. Acquiring MCS data with an 8-km-long streamer, as current *Langseth* capabilities allow, will further improve imaging of lower crustal and uppermost mantle features, as well as enabling amplitude-vs-offset studies of the physical properties of such deep targets. An added value of long-offset MCS data is their potential for conducting high-resolution waveform tomography studies to obtain the fine-scale seismic velocity structure of the upper- and mid-crust in the vicinity of a deep drill hole, thus contributing to a better integration of drilling sampling/results and regional geophysical data.

(3) 3D large-scale Ocean Bottom Seismometer (OBS) data. *Langseth* offers a superb powerful airgun array for active-source 3D wide-angle refraction/reflection OBS experiments. Using traveltime tomography techniques, these type of data allow resolving the 3D *P*- and *S*-wave velocity and anisotropy structure of the crust and uppermost mantle at scales of several kilometers, which will help interpretation of drilling results and placing them in the appropriate tectonic context.



(4) 2D high-resolution OBS data. Data acquired with a large number of densely spaced OBSs, in conjunction with *Langseth's* excellent seismic source, can be used for high-resolution determination of the velocity structure of the Moho transition zone (MTZ) in the vicinity of a drill hole using waveform tomography approaches [e.g., *Operto et al.*, 2006]. At long source-receiver offsets (~15-40 km), the MTZ seismic signature is a high-amplitude wide-angle reflection ideal for frequency-domain waveform tomography studies [e.g., *Brenders and Pratt*, 2007].

## References.

- Brenders, A. J., and R. G. Pratt (2007), Full waveform tomography for lithospheric imaging: results from a blind test in a realistic crustal model, *Geophys. J. Int.*, *168*, 133-151.
- Canales, J. P., M. R. Nedimović, G. M. Kent, S. M. Carbotte, and R. S. Detrick (2009), Seismic reflection images of a near-axis melt sill within the lower crust at the Juan de Fuca Ridge, *Nature*, *460*, 89-93.
- Kent, G. M., I. I. Kim, A. J. Harding, R. S. Detrick, and J. A. Orcutt (1996), Suppression of sea-floor-scattered energy using a dip-moveout approach; application to the mid-ocean ridge environment, *Geophysics*, *61*, 821-834.
- Nedimović, M. R., D. R. Bohnenstiehl, S. M. Carbotte, J. P. Canales, and R. P. Dziak (2009), Faulting and hydration of the Juan de Fuca Plate system, *Earth Planet. Sci. Lett.*, *284*, 94-102.
- Nedimović, M. R., S. M. Carbotte, A. J. Harding, R. S. Detrick, J. P. Canales, J. B. Diebold, G. M. Kent, M. Tischer, and J. Babcock (2005), Frozen magma lenses below the oceanic crust, *Nature*, 1149-1152.
- Operto, S., J. Virieux, J.-X. Dessa, and G. Pascal (2006), Crustal seismic imaging from multifold ocean bottom seismometer data by frequency domain full waveform tomography: Application to the eastern Nankai trough, *J. Geophys. Res.*, *111*, B09306, doi:09310.01029/02005JB003835.

### Appendix 3 - INVEST Meeting White Papers

A number of white papers related to drilling the oceanic lithosphere have been submitted to The IODP INVEST meeting (*IODP New Ventures in Exploring Scientific targets*, September 23-25, 2009, Bremen, Germany; <http://www.marum.de/iodp-invest.html>). Several of them directly or indirectly addressed scientific objectives and/or engineering/technological aspects of the MoHole project. They can be downloaded from the INVEST meeting web site :

- *Petrological nature of the oceanic Moho*, by Shoji Arai (Kanazawa University, Japan) : [http://www.marum.de/Binaries/Binary42200/Arai\\_MoHole.pdf](http://www.marum.de/Binaries/Binary42200/Arai_MoHole.pdf)
- *Integration of drilling into deep oceanic crust and seafloor geophysical observations for investigation of upper mantle structure and plate aging*, by Kiyoshi Baba (Earthquake Research Institute, University of Tokyo, Japan) et al. : [http://www.marum.de/Binaries/Binary42202/Baba\\_CrustMantleStructureIntergration.pdf](http://www.marum.de/Binaries/Binary42202/Baba_CrustMantleStructureIntergration.pdf)
- *Ocean Drilling and Exploring a Heterogeneous Ocean Crust*, by Henry J.B. Dick (Dept. of Geology & Geophysics, WHOI, Woods Hole, USA) et al. : [http://www.marum.de/Binaries/Binary42446/Dick\\_CrustalHeterogeneity.pdf](http://www.marum.de/Binaries/Binary42446/Dick_CrustalHeterogeneity.pdf)
- *Towards coordination and integration of deep marine biosphere research: the Dark Energy Biosphere Institute (DEBI)*, by Katrina J. Edwards (USC, Los Angeles, USA) et al. : [http://www.marum.de/Binaries/Binary42209/Edwards\\_DeepBiosphereDEBI.pdf](http://www.marum.de/Binaries/Binary42209/Edwards_DeepBiosphereDEBI.pdf)
- *Technological challenges necessary for the new horizon of IODP*, by Hiromi Fujimoto (Graduate School of Science, Tohoku University, Japan) : [http://www.marum.de/Binaries/Binary42212/Fujimoto\\_TechnologicalChallenges.pdf](http://www.marum.de/Binaries/Binary42212/Fujimoto_TechnologicalChallenges.pdf)
- *Vertical Magnetization Structure of Ocean Crust and uppermost Mantle*, by Toshiya Fujiwara (IFREE, JAMSTEC, Japan) : [http://www.marum.de/Binaries/Binary42214/Fujiwara\\_OceanCrustMagneticStructure.pdf](http://www.marum.de/Binaries/Binary42214/Fujiwara_OceanCrustMagneticStructure.pdf)
- *Evolution of Hydrothermal Circulation*, by Robert N. Harris (College of Oceanic and Atmospheric Sciences, Oregon State University, USA) et al. : [http://www.marum.de/Binaries/Binary42217/Harris\\_HydrothermalCirculationEvolution.pdf](http://www.marum.de/Binaries/Binary42217/Harris_HydrothermalCirculationEvolution.pdf)
- *Drilling deep through the ocean crust into the upper mantle*, by Benoît Ildefonse (Géosciences Montpellier, CNRS, Université Montpellier 2, France) et al. : [http://www.marum.de/Binaries/Binary42226/Ildefonse\\_MoHole.pdf](http://www.marum.de/Binaries/Binary42226/Ildefonse_MoHole.pdf)
- *Technological Drivers for Future IODP Science*, by the IODP Engineering Development Panel : [http://www.marum.de/Binaries/Binary42467/IODPEngineeringDevelopmentPanel\\_TechnologicalDrivers.pdf](http://www.marum.de/Binaries/Binary42467/IODPEngineeringDevelopmentPanel_TechnologicalDrivers.pdf)
- *New World of Technology developed with "Chikyu"*, by Yoshio Isozaki (CDEX, JAMSTEC, Japan) et al. : [http://www.marum.de/Binaries/Binary42443/Isozaki\\_NewTechnologyChikyu.pdf](http://www.marum.de/Binaries/Binary42443/Isozaki_NewTechnologyChikyu.pdf)
- *Technology Developments for IODP Phase 2*, by Japan Domestic INVEST Workshop Technology Development Group : [http://www.marum.de/Binaries/Binary42439/JapanDomesticINVESTWorkshopTechnologyDevelopmentGroup\\_Technology.pdf](http://www.marum.de/Binaries/Binary42439/JapanDomesticINVESTWorkshopTechnologyDevelopmentGroup_Technology.pdf)

- *Development of high temperature drilling technologies for 21st century Mohole*, by Engineering Development Advisory Committee (J-DESC, Japan) :  
[http://www.marum.de/Binaries/Binary42316/JapaneseEDGroup\\_HighTemperatureDrilling.pdf](http://www.marum.de/Binaries/Binary42316/JapaneseEDGroup_HighTemperatureDrilling.pdf)
- *Technological development for high temperature measurement in IODP*, by Engineering Development Advisory Committee (J-DESC, Japan) :  
[http://www.marum.de/Binaries/Binary42317/JapaneseEDGroup\\_HighTemperatureMeasurements.pdf](http://www.marum.de/Binaries/Binary42317/JapaneseEDGroup_HighTemperatureMeasurements.pdf)
- *Ultra deep water and ultra deep drilling technologies for 21st Century Mohole*, by Engineering Development Advisory Committee (J-DESC, Japan) :  
[http://www.marum.de/Binaries/Binary42283/Watanabe\\_UltradeepDrilling.pdf](http://www.marum.de/Binaries/Binary42283/Watanabe_UltradeepDrilling.pdf)
- *Geophysical studies toward deep drilling through the oceanic crust into the upper mantle*, by Shuichi Kodaira (IFREE, JAMSTEC, Japan) et al. :  
[http://www.marum.de/Binaries/Binary42243/Kodaira\\_GeophysicsUpperMantle.pdf](http://www.marum.de/Binaries/Binary42243/Kodaira_GeophysicsUpperMantle.pdf)
- *Benefits of in-situ and well-side geochemical monitoring at riser drilling*, by Hidenori Kumagai (IFREE, JAMSTEC, Japan) et al. :  
[http://www.marum.de/Binaries/Binary42244/Kumagai\\_GeochemicalMonitoringRiser.pdf](http://www.marum.de/Binaries/Binary42244/Kumagai_GeochemicalMonitoringRiser.pdf)
- *MOHOLE proposal for understanding the role of water in magmatic processes of oceanic ridges*, by Hiroaki Sato (Dept. Earth and Planetary Sciences, Kobe University, Japan) :  
[http://www.marum.de/Binaries/Binary42262/Sato\\_MoHole.pdf](http://www.marum.de/Binaries/Binary42262/Sato_MoHole.pdf)
- *Realistic Mohole using D/V Chikyu*, By Nobukazu Seama (Kobe University, Japan) et al. :  
[http://www.marum.de/Binaries/Binary42463/Seama\\_Mohole.pdf](http://www.marum.de/Binaries/Binary42463/Seama_Mohole.pdf)
- *The Ocean Lithosphere: A Fundamental Component of the Earth System (encapsulating discussions of the InterRIDGE-IODP "Melting, Magma, Fluids and Life" Workshop)*, by Damon Teagle (NOC, University of Southampton, UK) et al. :  
[http://www.marum.de/Binaries/Binary42445/Teagle\\_MeltingMagmaFluidLife.pdf](http://www.marum.de/Binaries/Binary42445/Teagle_MeltingMagmaFluidLife.pdf)
- *21st Century MoHole - Researches on Magmatic Processes of the Moho Transition Zone beneath the fast-spreading ridges*, by Susumu Umino (Kanazawa University, Japan) :  
[http://www.marum.de/Binaries/Binary42269/Susumu\\_MoHole.pdf](http://www.marum.de/Binaries/Binary42269/Susumu_MoHole.pdf)
- *Breakthrough the Discontinuity: 21st Century Mohole*, by Susumu Umino (Kanazawa University, Japan) et al. :  
[http://www.marum.de/Binaries/Binary42280/Umino\\_JapanWG\\_MoHole.pdf](http://www.marum.de/Binaries/Binary42280/Umino_JapanWG_MoHole.pdf)
- *Life and Ecosystem in Deep Biosphere and Subseafloor Aquifers*, by Hiroyuki Yamamoto (JAMSTEC, Japan) et al. :  
[http://www.marum.de/Binaries/Binary42307/Yamamoto\\_DeepBiosphere2.pdf](http://www.marum.de/Binaries/Binary42307/Yamamoto_DeepBiosphere2.pdf)

## Appendix 4 - Links to Earlier Scientific Planning Documents

- COMPLEX, 1999 conference report :  
[http://www.iodp.org/index.php?option=com\\_docman&task=doc\\_download&gid=808](http://www.iodp.org/index.php?option=com_docman&task=doc_download&gid=808)
- "Opportunities in Geochemistry for Post-2003 Ocean Drilling", 2000 workshop report :  
[http://www.iodp.org/index.php?option=com\\_docman&task=doc\\_download&gid=784](http://www.iodp.org/index.php?option=com_docman&task=doc_download&gid=784)
- ODP Architecture of the lithosphere PPG, summary report, 2003 :  
[http://www.iodp.org/index.php?option=com\\_docman&task=doc\\_download&gid=783](http://www.iodp.org/index.php?option=com_docman&task=doc_download&gid=783)
- Mission Moho, 2006 Workshop Report :  
[http://www.iodp.org/index.php?option=com\\_docman&task=doc\\_download&gid=1309](http://www.iodp.org/index.php?option=com_docman&task=doc_download&gid=1309)
- "Mission Moho: Formation and Evolution of Oceanic Lithosphere", EOS, 2006 :  
[http://www.iodp.org/index.php?option=com\\_docman&task=doc\\_download&gid=1302](http://www.iodp.org/index.php?option=com_docman&task=doc_download&gid=1302)
- "Mission Moho Workshop: Drilling Through the Oceanic Crust to the Mantle", Scientific Drilling, 2007 :  
[http://www.iodp.org/iodp\\_journals/2\\_Mission\\_Moho\\_Workshop\\_SD4.pdf](http://www.iodp.org/iodp_journals/2_Mission_Moho_Workshop_SD4.pdf)
- MissionMoho IODP proposal, 2007 :  
[http://www.gm.univ-montp2.fr/spip/IMG/pdf/MissionMohoProposal\\_April07\\_LowRes.pdf](http://www.gm.univ-montp2.fr/spip/IMG/pdf/MissionMohoProposal_April07_LowRes.pdf)
- Melting, Magma, Fluids and Life, 2009 workshop report :  
available at <http://www.interridge.org/WG/DeepEarthSampling/workshop2009>

## Appendix 5 - Workshop Presentation slides

### Introduction, scientific rationale for the MoHole

- **Damon A.H. Teagle & Mission Moho Proponents** - Beyond the Moho: Scientific justification for a complete penetration of intact in situ ocean crust and into the upper mantle ..... 65

### Available Site survey data in the Pacific

- **Douglas S. Wilson** - The Trade-off of Depth and Temperature in choosing a MoHole Site ..... 75
- **Nobukazu Seama** - Hawaii site in comparison with IODP site 1256 ..... 83
- **Douglas S. Wilson, Alistair J. Harding, Eric Hallenborg, & Graham Kent** - Site Survey Data at ODP Site 1256 (ODP Leg 206, IODP Expeditions 309/312) ..... 106
- **Donna K. Blackman** - “Deep Tow” Site: ~30Myr Old Crust off Southern California ... 115
- **Shuichi Kodaira** - IFREE seismic studies toward the Mohole project: results and future plan ..... 120

### Current understanding of technological feasibility

- **Greg Myers** - MoHole Engineering... Seeking solutions inside and outside of IODP .... 134
- **Shomei Kobayashi** - “Chikyu” Drilling Challenge for the MoHole ..... 144
- **Masaoki Yamao** - The CHIKYU Engineering Development ..... 157
- **Yuichi Shinmoto** - Technical Development of Coring System in the Quest to reach the Earth’s Mantle ..... 163

### Drilling and coring at high temperatures

- **David Mainprice & Philippe Pezard** - Drilling DEEP into HOT Oceanic Crust, Constraints and Challenges ..... 167
- **Alister Skinner** - Iceland Deep Drilling Project ..... 177

**Brief!**

## **Beyond the Moho: Scientific justification for a complete penetration of intact *in situ* ocean crust and into the upper mantle**

**Damon A.H. Teagle**  
on behalf of the Mission Moho Proponents

**Benoît Ildefonse** Natsue Abe, Peter B. Kelemen, Hidenori Kumagai,  
Damon A.H. Teagle, Douglas S. Wilson, Gary Acton, Jeffrey C. Alt,  
Wolfgang Bach, Neil R. Banerjee, Mathilde Cannat, Richard L. Carlson,  
David M. Christie, Rosalind M. Coggon, Laurence Coogan, Robert Detrick,  
Henry J.B. Dick, Jeffrey S. Gee, Kathryn Gillis, Alistair Harding, Jeffrey A.  
Karson, Shuichi Kodaira, Juergen Koepke, John MacLennan, Jinichiro Maeda,  
Christopher J. MacLeod, D. Jay Miller, Sumio Miyashita, James H. Natland,  
Toshio Nozaka, Mladen Nedimovic, Yasuhiko Ohara, Kyoko Okino, Philippe  
Pezard, Eiichi Takazawa, Takeshi Tsuji, Susumu Umino

## **Mission Moho**

**Ultimate Goal:**


**A complete section through intact ocean crust  
formed at a fast spreading rate  
into peridotites of the upper mantle**



*No longer Science Fiction!  
Now within the realm of  
technical feasibility*




## From the safe distance of the planetary scale.....



From Umino et al., INVEST 2009

## Project "Mohole" 1957 - 1966 American Miscellaneous Society



Offshore Guadalupe Island  
(Baja California, Mexico)  
**March-April 1961**

First deep water drillship  
Dynamic positioning  
Borehole re-entry  
~ 3500 mbsl

Drilled 5 holes to max depth 183 m,  
miocene sediments & ~ 14 m of basalt  
First sampling of Seismic Layer 2



## Project "Mohole" 1957 - 1966



### High Drama of Bold Thrust through Ocean Floor



JOHN STEINBECK AND PHOTOGRAPHER GORD ON DECK OF THE CUSS I

#### EARTH'S SECOND LAYER IS TAPPED IN PRELUDE TO MOHOLE

Last week Project Mohole (LIFE, April 7) made scientific history when its drilling barge, CUSS I (whose name is made up of the initials of oil companies who developed it: Continental, Union, Gulf and Republic), pierced 881 feet into the sea floor to get some samples of the earth's never-before-penetrated second layer. On board to describe the extraordinary operation for LIFE was novelist John Steinbeck, who is also an amateur oceanographer.

by JOHN STEINBECK



*John Steinbeck  
Life  
April 14, 1961*

*"Our expedition should destroy the old and well loved error that doers and thinkers are different breeds"*

Inside James Wood, Michel Faber, Barbara Ehrenreich, Mark Lawson, Charlie Brooker

# The Guardian

The joy of being 60 Michel Houellebecq

The rich, fit, frisky generation Weekend

The dark genius of HP Lovecraft Review

Every match, every player - free guide to the Lions' tour with Welchart

Inside Sport

Man to be charged with McCartney murder today

## Journey to the centre of Earth

### The search for life six miles down

Police investigate religious links after witchcraft abuse of child, 8

Harry Potter and the missing gun

## Project MoHole back in the News

New York Times  
The Sunday Times  
Newsweek  
Metro



Big George's Breakfast  
Daily Mail



Richard and Judy!



## Unfortunately...


Many of the major science motivations of  
Project MoHole and  
40 years of scientific ocean drilling  
still remain:

- Nature of the **Moho**
- **Seismic Layering** of the crust
- **Accretion** of the ocean crust
- Nature of **Magma Chambers**
- **Thermal Structure** of the ocean crust
- Geometry of **Hydrothermal Cooling**
- Chemical, fluid and isotopic **Fluxes**
- Ridge Axis vs **Ridge Flank Exchange**
- Origin of **Magnetic Stripes**

### Primary inhibitor

**Dearth of continuous sections of ocean crust**

### Basement drilling 1968 to 2005



Until recently:

1 penetration of lava-dikes - Hole 504B  
no sampling of dikes-gabbros

**Drilling Hard Rocks in the Oceans is Hard!**

## Mission Moho

### Ultimate Goal:

**A complete section through intact ocean crust  
formed at a fast spreading rate  
into peridotites of the upper mantle**


### Scientific objectives:

- Geological nature of the **Mohorovicic Seismic Discontinuity**
- ***In situ* composition** and physical properties of the **Uppermost Mantle**
- Upper **Mantle Dynamics** and **Melt Migration** processes
- **Bulk Composition** of the oceanic crust
- **Magmatic accretion** at mid-ocean ridges
- **Extent, location, and intensity of hydrothermal exchanges**
  - global chemical budgets, seawater chemistry, subduction fluxes
- Calibrate regional **Seismic Layering** against borehole and samples
- **Marine magnetic anomalies**
- **Limits of Life** in the Oceanic Lithosphere

### Ocean crust architecture is variable

Fast-spreading : East Pacific Rise

Slow-spreading : Mid-Atlantic Ridge




After Canales et al. (2003)

After Cannat et al. (1995)



## Why fast spread crust?




- **Relatively uniform**
  - Layered and relatively homogeneous
- **20% of modern ridges fast spreading**
  - 50% of modern ocean crust
  - ~30% of Earth's surface
  - Majority of crust subducted into mantle in past 200 Ma
- **Theoretical models for crustal accretion**
- **Oman ophiolite**
  - potential analog?

Slow spread crust: heterogeneous – characterization beyond the scope of MoHole

### **BUT – Tectonic Windows: Slow and Fast Spread Crust:**

Near term science achievements

**Opportunities to learn how to drill deeply in oceanic basement?**



## What is the Moho?


Primary acoustic interface within the Earth ( $V_p > 8$  km/s)

Fundamental to geodynamics of our planet

Oceans ~5-8 km, Commonly a bright reflector

**Seismic Discontinuity ASSUMED to be**

**boundary between Crust and Mantle**



Is the Moho:

- mafic cumulates & residual peridotites?
- mafic & ultramafic cumulates?
- serpentinitized peridotites & peridotites?
- gabbro-intruded peridotites & peridotites?

How sharp is the boundary?

Is the Moho different in different places?

Dick, Natland and Ildefonse, 2006

## In Situ Samples and Observations of the Convecting Upper Mantle

- **Chemical Composition and Physical Properties of MORB source**

$f_{O_2}$ , S, CO<sub>2</sub>, CH<sub>4</sub>, Graphite, H<sub>2</sub>O, Li, B, He, noble gases

U, Th - grain boundaries

- Scales of **Mantle Source Heterogeneity** - Sr, Nd, Pb, Hf, Os
- Grain size and **Deformation** history - seismics, melt transport, corner-flow
- **Melt Focusing and Extraction**




## In Situ Samples and Observations of the Convecting Upper Mantle

- **Chemical Composition and Physical Properties of MORB source**

$f_{O_2}$ , S, CO<sub>2</sub>, CH<sub>4</sub>, Graphite, H<sub>2</sub>O, Li, B, He, noble gases

U, Th - grain boundaries

- Scales of **Mantle Source Heterogeneity** - Sr, Nd, Pb, Hf, Os
- Grain size and **Deformation** history - seismics, melt transport, corner-flow
- **Melt Focusing and Extraction**



For Buoyancy-driven flow


$V_{\text{uppermantle}} > V_{\text{plate}}$

Inversion of shear sense  
close to Moho

(e.g., Oman, Nicolas et al., 1994; Ildefonse et al., 1995)




## What is the coupling between plate and mantle motions?



Harmon et al., 2004, GRL

## Geophysical evidence for mid-ocean ridge magma chambers



### Seismic experiments at modern MOR

#### Multi-channel reflection:

- bright axial reflector
- 1-3 km below seafloor
- 100s m wide x 10s m thick
- partially (usually) to fully molten
- **interpreted** to be magma chamber


#### Refraction velocity mapping:

- Low-velocity zone beneath AMC reflector
- LVZ extend to base of crust
- low melt fraction 1-2%
- transmits s-waves
- Solid, full-thickness crust within 2-3 km from axis

## Test models of Crustal Accretion

### Gabbro-Glacier


Henstock et al., 1993,  
Phipps-Morgan and Chen, 1993,  
Quick and Denlinger, 1993



Conductively Cooled  
Lower crust

### Sheeted Sills


Boudier et al., 1996  
Kelemen et al., 1997  
MacLeod and Yaouancq, 2000



Hydrothermally Cooled  
Lower Crust

## Conductive or convective cooling?

### Predicted profiles that can be tested with drill cores



Original fig from Roz Coggon

## Criteria for Siting the MoHole

Heavy Metal  
vs  
Hard Rock



We need to learn  
how to drill  
deep hot rocks

- Water depth within **Riser Capability** <4000 m (4500 m ?)
- Age >15 Ma, preferably >20 Ma, Mantle T <200°C
- **Weather window** >8 months, calm seas, gentle currents
- **Formation at fastest possible spreading rate** (>80 mm/yr)
- **Continuous layered structure**, low abyssal hills
- **Simple tectonic setting** (avoid seamounts, fracture zones)
- **Well Imaged Moho** (reflection and refraction)
- **Original latitude** >±15° - marine magnetic anomalies
- **Sediment thickness** to support riser hardware (>50 m)
- **Close to major port facilities**
- **Slightly below average crustal thickness**

**No Site fulfils ALL Criteria**

# The Trade-off of Depth and Temperature in choosing a MoHole Site


Doug Wilson, UCSB

## Desired Site Properties:

- Not too deep ( $< \sim 4$  km: riser, pipe weight, etc.)
- Not too hot at Moho ( $< \sim 200^\circ\text{C}$ )
- Nearly normal ocean crust
- Need to settle for 2 out of 3




## Observed Sea Floor Depth vs Age




Most sites subside deeper than 4000 m by 25 Ma

## Predicted Temperature as a Function of Age and Depth




At 6 km, cooling below  $200^\circ\text{C}$  occurs after 25 Ma

## Temp. vs. Water Depth, Average Crust



Depth from Sclater curve, temperature from half-space model

## Temp. vs. Water Depth, Average Crust



Depth from Sclater curve, temperature from half-space model



## High-Priority Site Attributes

- Age  $>\sim 15$  Ma
- Depth  $<\sim 4$  km
- Intermediate to fast spreading rate, abyssal hill faults  $< 200$  m high
- Weather window 9-12 months
- Simple, mapped tectonics


## High-Priority Site Attributes

- Age  $>\sim 15$  Ma
- Depth  $<\sim 4$  km
- Eastern Pacific,  $35^{\circ}\text{N}$ – $35^{\circ}\text{S}$
- Simple, mapped tectonics


## Desirable Site Attributes

- Age > 20 Ma
- Close to major port(s)
- International waters or member EEZ
- Crustal thickness 5.0–5.5 km
- 12-month weather window
- Original latitude >  $\pm 15^\circ$
- Fastest available spreading rate


Depth  
limited by  
4-km riser



Age > 15  
Ma




Depth < 4 km  
Age > 15 Ma  
Simple & well  
mapped




## Potential MoHole Sites

Depth < 4 km




Depth < 4.5 km




## Potential MoHole Sites

Depth < 4 km




Depth < 3.5 km





## Temp. vs. Water Depth, Average Crust



Depth from Sclater curve, temperature from half-space model

## Concluding Remarks

- Many potential sites are available for ~30 Ma at water depths 4000-4300 m, probable Moho temperature ~180°C.
- A few potential sites are available for 15-20 Ma at water depths 3400-3700 m, probable Moho temperature ~200-250°C.
- Which is a bigger engineering challenge: 4200 m water depth or 240°C formation?

# Hawaii site in comparison with IODP site 1256

Nobukazu Seama  
Kobe University

## How should we decide a Mohole site?

- Recent high quality seismic surveys in the western Pacific using air-gun, multi-channel streamers, and ocean-bottom seismometers, indicate that velocity structure of ocean crust and of uppermost mantle has **variety** even if it was formed at a fast spreading rate ridge system.
- The Mohole drilling provides **a deep reference hole** for the ocean crust and the uppermost mantle.
  - to properly interpret the velocity structure
  - to point out the similarity and/or the difference in velocity structure in space (or crustal age)




## How should we decide a Mohole site?

- Detailed seismic velocity structure and its characters below are especially important.
  - Crustal velocity structure (velocity value, etc.)
  - Moho discontinuity (amplitude of reflector; single or multiple reflectors, etc.)
  - Uppermost mantle velocity structure (velocity value and degree of anisotropy, etc.)
- The final Mohole site should be decided after discussions among the international community with enough geophysical data for all candidates.

If technology allows us to drill  
wherever we like,

Then, we discuss where show best seismic velocity structure as a deep reference hole for the ocean crust and the uppermost mantle.

- Technology limitation forces us to discuss feasible drilling sites with near future technology, where we should do high quality seismic surveys in a few years.




## Site 1256

In Mission Moho WS full report,


- Current primary site
- The greatest currently perceived threat at Site 1256 is from potentially high down-hole temperatures. For this reason, the alternate site should be on older (colder) crust.

→ Bench mark for the alternate sites


# Temperature in 1256D




## Temperature estimation



## Temperature estimation




## Temperature estimation






# Temperature estimation




## Site 1256 summary


- For the coming next drilling, temperature measurements are essential before and after the drilling.
- The alternate sites should be on older crust, because potentially high down-hole temperatures in Site 1256.
- Site 1256 is a bench mark for the alternate sites




Off Mexico








Load of Hawaii island chains bends  
the lithosphere



# Lithosphere bending



JOURNAL OF GEOPHYSICAL RESEARCH

VOL. 69, No. 8

APRIL 15, 1964

## Mohole Site Selection Studies North of Maui<sup>1</sup>

GEORGE G. SHOR, JR., AND DWIGHT D. POLLARD<sup>2</sup>



*Scripps Institution of Oceanography, University of California, San Diego*

**Abstract.** Seismic refraction profiles made across the Hawaiian arch, the Hawaiian deep, and onto the Hawaiian ridge north of the island of Maui indicate that the Mohorovicic discontinuity lies at a depth of about 10.5 km beneath the arch and dips to 13 km beneath the deep and the ridge. The variation in depth to the mantle from arch to deep is caused principally by variations in the thickness of the sediments and volcanics; the material having velocities between 5 and 7 km/sec has almost constant thickness. The hypothesis of Vening Meinesz that the Hawaiian deep is caused by down-warping of the crust under the load of the Hawaiian ridge, and that the Hawaiian arch is a side bulge caused by elastic yielding, is substantiated by the data obtained off Maui. Menard's hypothesis that the smooth sea floor on the Hawaiian arch is caused by extensive lava flows rather than by sedimentation is also supported; material having a velocity near 4 km/sec at or just beneath the sea floor was found under the deep and under the south side of the arch. This material has apparently filled the original deep in part. Some evidence exists for lateral variation of the velocity of compressional waves in the mantle. This, plus variations of velocity in the shallower layers and possible changes of dip of layers within the length of individual refraction profiles, makes the computed depth to mantle at individual stations depend somewhat on the form of solution and pairing of stations. Depths to the discontinuity as shallow as 9 km may exist at some points on the arch. A small area in which high-velocity material occurs within 7 km of sea level was found under the shelf northwest of Kahului, Maui; this may be material brought up from the mantle either in a volcanic pipe or by faulting.




## Hawaii site

- Crustal Age:  
80Ma
- Half spreading rate:  
35-40 mm/yr
- Original latitude:  
near equator
- Water depth:  
4050-4300 m



## Previous seismic survey results


- Clear Moho discontinuity



Watts et al., 1985

## Previous seismic survey results


- Clear Moho discontinuity




## Previous seismic survey results

- Clear Moho discontinuity

ten Brink and Brocher, 1988




Lindwall, 1991





## Geomagnetic Anomaly

- Clear 2D Normal/Reverse patterns
- Crustal age is well constrained ~80Ma
- Half spreading rate 35-40 km/Myr




## R/V Kilo Moana KM1003 Cruise



B. Taylor, G. Moore, G. Ito, 1 post doc,  
6 students (SOEST, Univ. Hawaii), N.  
Abe, T. Fujiwara (IFREE, JAMSTEC)

- 20 Feb ~ 25 Feb 2010 (6 days)





## KM1003 with R/V Kilo Moana

- Swath bathymetry (Simrad EM120)
- Sub-bottom profiler (Knudsen 320)
- Gravity (LaCoste-Romberg S-33)
- Magnetics (Geometrics G-882 Cesium)




Gravitymeter





Magnetometer

After Fujiwara








# Could Hawaiian Plume modify the crust and uppermost mantle?


- 1) Infill material
- 2) Arch Volcanoes



Wolfe et al., 1994

## Previous seismic survey results

- Deep crustal sill complex




Watts et al., 1985



# Previous seismic survey results


- Deep crustal sill complex



Watts et al., 1985


## Could Hawaiian Plume modify the crust and uppermost mantle?

- 1) Infill material
- 2) Arch Volcanoes




Bianco et al., 2005

# Arch Volcanic Fields





Bianco et al., 2005

# Arch Volcanoes




Normark  
et al., 1989






## Seismicity could be indicator for magma reservoirs and conduits



Klein et al., 1987


## Hypocenters






# Deep S structure


Wolfe et al., 2010




# Heat Flow



Stein and Stein, 1992




## Proposed seismic survey lines



## Temperature estimation for Hawaii site

Half space thermal model + Square root  $t$  low  
 → slightly higher than  $100^{\circ}\text{C}$  at  $6\text{km}$  →  $<150^{\circ}\text{C}$

### Predicted Temperature as a Function of Age and Depth After Wilson



At 6 km, cooling below  $200^{\circ}\text{C}$  occurs after 25 Ma

Good for

- Drilling
- LWD

(Logging-While-Drilling)

- Microbiology

(Temperature for amino acid surviving)

×4




## Hawaii site shows desirable site attributes?


- Nearly normal ocean crust → Need to avoid Arch volcanoes
- Age > 20Ma ← Not too hot at Moho (<~200°C)  
○ <150°C
- Not too deep (4-km riser) → 4050-4300 m
- Close to major port(s) ○
- International waters or member EEZ ○
- Crustal thickness 5.0-5.5 km → ~6 km?
- 12-month weather window ○
- Original latitude > ±15 → nearly equator
- High quality seismic survey → We need to do it
- Slower spreading rate than current fastest ridge system

## Desirable item: Slower spreading rate than current fastest ridge system

<~80mm/yr half spreading rate




Mohole (reference hole)



## Weak points of Site 1256

- Too hot at Moho?
- Faster spreading rate than current fastest ridge system ( $\sim 110\text{mm/yr}$  half spreading rate)
- Clear Moho?  $\rightarrow$  High quality seismic survey




## How should we decide a Mohole site?

- $\rightarrow$  Detailed seismic velocity structure and its characters below are especially important.
  - Crustal velocity structure (velocity value, etc.)
  - Moho discontinuity (amplitude of reflector; single or multiple reflectors, etc.)
  - Uppermost mantle velocity structure (velocity value and degree of anisotropy, etc.)
- $\rightarrow$  The final Mohole site should be decided after discussions among the international community with enough geophysical data for all candidates.


# Site Survey Data at ODP Site 1256

## ODP Leg 206, IODP Expeditions 309/312


Douglas S. Wilson, Alistair J. Harding, Eric Hallenborg, Graham Kent





## EW9903 Trackline





## Isochrons and site-survey track









## Seismic reflection section











Line 21 original 1-D  
velocity model

Site chosen at fairly high upper crust velocity of 4.9-5.0 km/s,  
Indicative of low porosity and hopefully, good drilling conditions







GUATB03: Line 22 Synthetic and Data at Line 25 Crossing




GUATB03: Line 21 at line 27 crossing






## Topography (or lack thereof) on Moho






## Comparison of refraction interfaces with reflection image




# “Deep Tow” Site: ~30 Myr Old Crust off Southern California



*Summary by Donna Blackman  
Scripps Institution of Oceanography*

## “Deep Tow” Location



*32°25'N 125°45'W, ~750 km offshore California*

*31-32 Myr old crust (magnetic anomaly & basalt date)*

*western flank of subducted East Pacific Rise*

*north of series of seamounts*



# What is Known About Site?

Luyendyk, Geological Society of America Bulletin, v 81, p2237, 1970

Deep-Tow geophysical package

sonar (sidescan & bathymetry)

3.5 kHz profiler

35 mm camera


magnetometer

airgun seismics (sediment profiling)

Coverage

2 Abyssal Hills and intervening trough

area ~30x20 km transected by several lines (dotted)



shipboard bathymetry (40-m contour interval)

## Local Geology of Site

EPR spreading rate during formation similar to R2K 9°N site (11 mm/yr, full rate)

Deep-tow data used to infer different origins for the 2 abyssal hills

eastern hill- fault-controlled horst block (overlying sediment reflector offsets)

western hill- elongate volcanic shield (morphology, basement reflectors)

Sediment

40-60 m thick

thin, continuous reflections- 10's cm thick volcanic ash (piston core & dredge)

post-deposition offsets, basement abutments overlain by unconformity >> faulting





Figure 6. Three-dimensional scale model of the bathymetry mapped in Figure 5. Each laminae represents 20 m. No vertical exaggeration. Bar at lower right of model is 2 km long. Model courtesy of G. Sheary, Bethlehem Steel Corporation.


model uses Deep-Tow high resolution bathymetry & illustrates different character of east/west abyssal hill



# Volcanic Knobs



acoustic records: 3.5 kHz profile (top), side-scan (lower)




local areas of little/no sediment  
disruption of sediment in at least 1 case  
sonar reflectivity

# Heat Flow

*Luyendyk concludes volcanic activity significantly post-dates initial formation of crust*

- $\sim 84 \text{ mW/m}^2$  predicted by Age model for 31 Myr crust
- values as high as  $370 \text{ mW/m}^2$  reported within 10 km (white dot)



# Inferred Structure

Luyendyk, 1970

*faulting & later volcanism begin ~10 m.y. ago (sedimentation rate, angular rock on sediment)*





Figure 24. A model of the formation of the abyssal hills in the study area. The western hill formed primarily from volcanic activity at the crest of the East Pacific Rise, and is probably representative of other hills that have been formed at the crest of a steadily spreading rise. The eastern hill has been created, and the western hill modified, by faulting of the oceanic crust long after the crust was formed. The sediment structure is a generalized cross section across the area. The upper section of the second layer is shown diagrammatically here as a complex of vertical dikes with a thin veneer of volcanic effusives on its surface. Vertical exaggeration is times 10; the sound velocity in the sediment is assumed to be 1.5 km/sec.

## Supporting Figures (Luyendyk, 1970)



Figure 20. Scattered angular rocks seen on the east slope of the volcanic hill (Fig. 11). The width of the field of view is between 8 and 9 m.

*interpreted as  
lava fragments*




*sediment abuts basement of eastern hill,  
is overlain unconformably*




# Other Studies in the Vicinity

*G. Shor refraction measurement, 70 km NW*


Layer	Velocity (km/s)	Thickness (km)	Depth (km)
0	1.502	4.28	—
1	2.15	0.37	4.28
2	5.88	0.82	4.85
3	6.82	4.62	5.47
4	7.91	—	10.09



*Seismic Reflection (NGDC, mostly single channel)*



*Seismic Refraction (USGS 1984, probably sonobuoy)*



## Remarks

*coverage of the Deep-Tow site is not up to modern-day standards but some worthwhile information is available:*

*documentation of local bathymetry and stratigraphy is sufficient and indication of seafloor character (photos) is OK*

*available heat flow and occurrence of basement knobs, inferred to be volcanic, raises serious concern*

*if Luyendyk estimate of activity, ~10 m.y. and perhaps continuing, is correct thermal structure may be hotter than typical for 30 Myr crust*

*a single measure of "typical" Moho depth, determined using constant-layer modeling (not gradients) that is 70 km away*

*no opportunity to place results into context of ridge evolution since spreading axis subducted*



# IFREE seismic studies toward the Mohole project: - results and future plan -




## Strategy and Plan

- 1<sup>st</sup> stage: ~ 2010
  - Seismic imaging of "typical" oceanic crust in the northwestern Pacific
  - Modeling Moho reflection based on ophiolite
- 2<sup>nd</sup> stage: 2011 ~ 2015
  - Integrated geophysical study in Mohole candidate areas
- 3<sup>rd</sup> stage: 201? ~ 202?
  - Geophysical studies while- and post-drilling
    - Core - log - seismic integration
    - Long-term monitoring (geophysical / geochemical)




# Seismic characters of oceanic crust, mantle and Moho in the NW Pacific

IAMSTEC NW-Pacific





JCG profiles






3

## JCG Marcus profiles





Oikawa et al. 2009
















## Northwestern Pacific


25 km

Moho becomes obscure toward trench

6000 m



## Mantle velocity variation




JAMSTEC

<http://www.jamstec.go.jp/>

12




## Mantle velocity variation




Velocity perturbation below basement derived from first arrival tomography



## Mantle velocity variation





Velocity perturbation below basement derived from first arrival tomography

















## Strategy and Plan

- 1<sup>st</sup> stage: ~ 2010
  - Seismic imaging of "typical" oceanic crust in the northwestern Pacific
  - Modeling Moho reflection based on ophiolite
- 2<sup>nd</sup> stage: 2011 ~ 2015
  - Integrated geophysical study at possible Mohole areas
- 3<sup>rd</sup> stage: 201? ~ 202?
  - Geophysical studies while- and post-drilling
    - Core - log - seismic integration
    - Long-term monitoring (geophysical / geochemical)



## Modeling Moho reflection based on ophiolite




Oman ophiolite

Figure courtesy of Prof. S. Arai



## Modeling Moho reflection based on ophiolite




Oman ophiolite

Figure courtesy of Prof. S. Arai





## Modeling Moho reflection based on ophiolite




Oman ophiolite


Figure courtesy of Prof. S. Arai

## Strategy and Plan

- 1<sup>st</sup> stage: ~ 2010
  - Seismic imaging of "typical" oceanic crust in the northwestern Pacific
  - Modeling Moho reflection based on ophiolite
- 2<sup>nd</sup> stage: 2011 ~ 2015
  - Integrated geophysical study in Mohole candidate areas
- 3<sup>rd</sup> stage: 201? ~ 202?
  - Geophysical studies while- and post-drilling
    - Core - log - seismic integration
    - Long-term monitoring (geophysical / geochemical)



## Need international cooperation







# MoHole Engineering...

## Seeking solutions inside and outside of IODP

MoHole Workshop  
June 3-5, 2010  
Kanazawa, Japan

Greg Myers  
Consortium for Ocean Leadership



## Outline

- The Challenge
- In the IODP toolbox
- Where do technology gaps exist
- RMR– A possible solution to borehole problems
- Summary



## The Challenge

- Water depth over 4km
- Hole depth over 6km - 7km
  - 10 Empire state buildings of water (4.5km)
  - 14 Empire State Buildings into the seafloor (6km)



## The Challenge

- Water depth over 4km
- Hole depth over 6km-7km
  - 10 Empire state buildings of water (4.5km)
  - 14 Empire State Buildings into the seafloor (6km)







## Drilling to MOHO is easy..all you have to do is...

- Stabilize the vessel to drill/core/log in water depths ~12,000 feet (3,600 meters) and greater
- Clean the borehole hole
- Keep the borehole vertical
- Manage pressure within the borehole
- Manage temperature within the borehole
- Manage stress within the borehole
- Collect samples, return all equipment
- Avoid unfavorable met-ocean conditions
- Find the funding and stay within time and financial constraints



## Site Selection Issues

- **Drilling:** Water depth, Moho depth, expected borehole temperatures, lithological variations, pressures, stresses, core size, etc
  - This will affect vessel parameters, pipe specifications, mud composition and volume, casing specs, etc
- **Logistics:** Resupply, crew rotations, permitting
  - This will affect, vessel parameters, mobilizations/demobilizations and access
- **Weather** (Met ocean): Average sea state, storm frequency, currents, commercial fishing, whale migrations
  - This affects vessel parameters, heave compensation, drive offs, crew preparation, dynamic positioning equipment need



## Critical Questions

- Can the MoHole science community deviate from status quo techniques for sampling? How about...
  - Subsea drilling systems
    - Presently, uses for shallow hole site characterization
    - Especially useful for upper 100m of young crust
    - Full depth systems under development (Seabed Rig AS)
  - Punch and go exploration
    - Drill, recovery all cuttings, wireline log, spot core or sidewall core
    - Lower cost, drill deeper
    - Latest generation logging tools needed
- What does the science community define as MoHole Success?
  - Sampling certain amount of crust, vs, upper mantle, coring, fluids, logging, etc
- How many holes must be drilled?
  - Can a multiple site program work? (1256D and Hawaii sites?)
  - Pilot hole with main hole
  - Multiple holes
  - Holes to be established for observatories?



## The IODP Toolbox: Building upon Scientific Ocean Drilling technological successes

- Mission specific platforms
- Chikyu and riser drilling
- JOIDES Resolution
- Observatories
- In-situ sampling of gabbros, hydrates, microbiological communities, overpressured sediments, any many more unique materials







## Technology Gaps: Deep Holes and Low Core Recovery

**Table T3.** Summary of coring bits used during Expedition 309.

Bit number	Manufacturer	Serial number	Bit type	Size	Number of cores	Interval cored (m)	Recovered (m)	Recovery (%)	Coring time (h)	Rate of penetration (m/h)
Log bit					Logging				43.8	
1256D-1	RBI	BF-739	CC-9	9 <sup>7</sup> / <sub>8</sub>	11	69.1	25.20	36.5	51.7	1.34
1256D-2	RBI	BF-852	CC-9	9 <sup>7</sup> / <sub>8</sub>	11	76.8	18.02	23.5	52.1	1.47
1256D-3	RBI	BF-854	CC-9	9 <sup>7</sup> / <sub>8</sub>	11	61.1	14.85	24.3	52.8	1.16
1256D-4	RBI	BF-856	CC-9	9 <sup>7</sup> / <sub>8</sub>	4	20.4	9.52	46.7	17.8	1.14
1256D-5	RBI	BF-858	CC-9	9 <sup>7</sup> / <sub>8</sub>	15	72.1	20.56	28.5	50.1	1.44
1256D-6	RBI	BF-741	CC-9	9 <sup>7</sup> / <sub>8</sub>	12	57.6	21.45	37.2	50.8	1.13
1256D-7	RBI	BF-742	CC-9	9 <sup>7</sup> / <sub>8</sub>	8	36.3	17.70	48.8	42.5	0.85
1256D-8	RBI	BF-853	CC-9	9 <sup>7</sup> / <sub>8</sub>	12	58.6	17.74	30.3	57.8	1.01
1256D-9	RBI	CL-540	CC-9	9 <sup>7</sup> / <sub>8</sub>	12	51.3	37.57	73.2	53.1	0.97
Log bit					Logging				76.0	
Totals:					96	503.3	182.61			1.17



## Technology Gaps: Deep Holes and Low Core Recovery

**Table T6.** Operations Summary, Expedition 312.

Hole	Latitude	Longitude	Number of cores	Interval cored (m)	Core recovered (m)	Recovery (%)	Drilled/washed (m)	Total penetration (m)	Time on hole (h)	Time on hole (days)
1256D-1	6°44.1631'N	91° 56.0612'W	0	0.00	0.00	0.0	0.00	0.00	61.50	2.6
1256D-T	6°44.1631'N	91° 56.0612'W	0	Drill ahead with 9-7/8 inch tricone bit to bottom					63.25	2.6
1256D-2	6°44.1631'N	91° 56.0612'W	11	54.60	8.58	15.7	0.00	54.60	99.75	4.2
1256D-3	6°44.1631'N	91° 56.0612'W	8	33.80	5.54	16.4	0.00	33.80	86.00	3.6
1256D-4	6°44.1631'N	91° 56.0612'W	6	24.00	1.31	5.5	0.00	24.00	78.25	3.3
1256D-5	6°44.1631'N	91° 56.0612'W	4	5.30	0.53	10.0	0.00	5.30	53.50	2.2
1256D-F1	6°44.1631'N	91° 56.0612'W	Fishing run 1 (9 inch fishing magnet with 2 junk baskets)						23.75	1.0
1256D-F2	6°44.1631'N	91° 56.0612'W	Fishing run 2 (9-1/2 inch fishing mill with 2 junk baskets)						28.00	1.2
1256D-F3	6°44.1631'N	91° 56.0612'W	Fishing run 3 (9-1/2 inch fishing mill with 1 junk basket)						28.50	1.2
1256D-F4	6°44.1631'N	91° 56.0612'W	Fishing run 4 (9 inch fishing magnet with 2 junk baskets)						22.05	0.9
1256D-6	6°44.1631'N	91° 56.0612'W	8	25.80	1.39	5.4	0.00	25.80	85.70	3.6
1256D-7	6°44.1631'N	91° 56.0612'W	12	46.00	10.68	23.2	0.00	46.00	96.00	4.0
1256D-8	6°44.1631'N	91° 56.0612'W	13	62.50	18.49	29.6	0.00	62.50	99.50	4.1





## Technology Gaps: Deep Holes and Low Core Recovery

**Table T1.** Hole summary, Expedition 315.

Hole	Location	Water depth (mbsl)	Number of cores	Drilled depth (m)	Interval cored (m)	Recovered (m)	Recovery (%)	Comments
C0001E	33°14.3442'N, 136°42.6924'E	2198.0	13 HPCS	118.10	118.10	112.67	95.4	Lost inner core barrel in hole
C0001F	33°14.3437'N, 136°42.7067'E	2197.0	19 HPCS, 2 ESCS	248.83	140.80	137.50	97.7	
C0001G	33°14.3237'N, 136°42.6933'E	2196.5	NA	74.50	NA	NA	NA	ROV cable tangled around drill pipe
C0001H	33°14.3233'N, 136°42.6840'E	2197.0	26 RCB	590.50	228.60	126.30	55.2	Hole caving
C0001I	33°14.2030'N, 136°42.4330'E	2198.5	NA	520.00	NA	NA	NA	Hole caving
Site C0001 totals:			32 HPCS, 2 ESCS, 26 RCB	1551.93	487.50	376.47	77.2	
C0002B	33°17.9928'N, 136°38.2029'E	1937.5	66 RCB	1057.00	582.00	208.30	35.8	
C0002C	33°18.0026'N, 136°38.1869'E	1936.6	2 HPCS	13.77	13.80	13.77	99.8	
C0002D	33°18.0075'N, 136°38.1910'E	1937.1	16 HPCS, 2 ESCS	204.00	204.00	161.90	79.4	
Site C0002 totals:			18 HPCS, 2 ESCS, 66 RCB	1274.77	799.80	383.97	48.0	



## Ultra-deep Drilling Statistics

Site	Water Depth (m)	Borehole Depth (m)	Total Depth (m)	Comments
Proposal 698Full-2	1,798	8,000	9,798	1 year? <250°C
KTB	0	9,101	9,101	4+ years; \$338 million; 265°C
Kola SG-3	0	12,262	12,262	24 years; 190°C
Bertha Rogers 1-27	0	9,583 (31,441ft)	9,583	1974 gas well
Nankai NT3-01	~2,000	6,000	8,000	450 days allocated; ~175°C
1256D	3,635	1,507	5,142	~5 months; ~70 °C
JR			10,290 (SODV)	Total string length
Deepest hole	3,463	2,111	5,574	190°C, Site 504B
Deepest water	5,980	560	6,540	
Chikyu (riser)	2,500 (max)	7,000 (max)	9,500	<250°C borehole
Deepest hole	500	3,700	4,200	Off Australia, non-IODP
Deepest water	2,200	2,700	4,900	
Chikyu (non-riser)		7,000 (max)	10,000	
Deepest hole	1,936	1,401.5	3,337.5	C2 Leg 314
Deepest water	4,081	494	4,575	

Table created by Bill Ussler - MBARI



## Needed downhole equipment



### Drilling and Coring

- Instrumented drill collars
- Microbio core barrels
- Transition zone corers
- High temperature mud and equipment
- Hyper-deepwater mud circulation



### Logging



- Drill collar fluid sampling and seismic measurements
- Latest generation wireline tools deployable from all platforms
- Fluid samplers, sidewall corers, geochemistry, magnetic resonance etc



## Borehole Management

- Managing the borehole means:
  - Remove cuttings
  - Provide lithostatic and pore pressure compensation
  - Develop mud cake on borehole wall to provide additional stability
  - Mitigate fluid inflows and outflows
  - Limiting excess pumping rates
- Historically, seawater with occasional mud sweeps has been utilized, thus the deepest IODP hole is 2,111m deep
- Engineered mud must be circulated continuously as part of a comprehensive plan to drill and core effectively









## Deepwater Riserless Mud Recovery Equipment



In-line pump



Mud suction module

Proposal received from AGR to design and build a 5,000 meter system



## Riserless Mud Recovery Timeline

- 2009 - Feasibility project – completed by IODP-MI
  - Demonstrated RMR feasibility for IODP to 3,650 m (on paper)
  - Funded by DeepStar Industry Consortium
- 2011-2012 Field Trial from an IODP platform in <3,650 m of water
  - Must be preceded by procurement of funding and completion of engineering, vessel modifications and operations simulation
  - Funded partially by DeepStar, RPSEA, major Industry operator/s, cost sharing by AGR and IODP operators
- 2013-2014 Field trial in water depth >3,650 m
  - Must be preceded by procurement of funding and completion of engineering, vessel modifications and operations simulation
- 2015 – Ultra-deephole in hyper-deepwater capability could be ready



## If we are to achieve IODP MOHO goals...



1. We cannot continue to rely so heavily on volunteer-based engineering. We have extremely high expectations for the technical panels, yet we must bolster our engineering resources from outside of IODP
2. Centralized, project-specific, engineering management with hard links to oil and gas are required to get the technical support, and perhaps financial support
3. We must make difficult decisions about the tradeoffs between continuous coring and punch and go drilling



## In Summary


- Drilling to MOHO appears technologically feasible
- IODP has been hugely successful and its coring and logging experience base must be tapped
- IODP does not have the requisite experience with drilling multiyear, multiphase ultra-deephole projects, thus we need contractual links with those who do
- The engineers cannot move forward confidently with feasible solutions, until we know what constitutes success (how deep, how much sampling, how hot, etc.)
- Lets make some progress together this week





eWS Kanazawa 10/Jun/03/





### Advantages of Riser drilling

	Riser	BOP	Mud Circulation
Well Control / Formation Pressure Control	✓	✓	✓
Borehole Stability	✓		✓
Deep Penetration	✓		✓
Large & Safe Wireline Logging tools run - passage	✓		✓
Large & Safe Casing run - passage	✓		✓
Better Core Recovery	✓		✓

## Roles of Drilling Mud

### Removing Cuttings

The mud circulated through the drill bit removes cuttings as they accumulate at the drill bit. These cuttings are circulated up to the ship through the annulus between the drill pipe and the borehole and the riser pipe.

### Stabilizing Borehole

The mud forms thin film (mud cakes) on the borehole walls and this helps prevent borehole collapse. By increasing the density of the mud, the hydrostatic pressure can be balanced with the pore pressure to prevent the influx from formation.

### Cooling & Lubrication of the Bit


The mud acts as a lubricant for the bit and protects the bit from wear and tear. It also cools the bit against the rising temperatures inside the Earth.




MoHoleWS Kanazawa 10/Jun/03/

## Roles of Casings

1. Protect Bore Hole Collapse
2. Separate communication between High/Low Pressure Zones
3. Basement for Wellhead System
4. Reduce Friction between Drill String and Borehole
5. Protect Borehole Observation tool




MoHoleWS Kanazawa 10/Jun/03/



## 2009 NanTro SEIZE Stage2

## NT2-11 Hole Drilling Sequence





## Limitless Wireline Logging Measurements

- ❖ Choice of combinations for hostile environments
- ❖ Dipole shear sonic, Fullbore electrical imaging, Dynamic formation testing and sampling, Magnetic resonance tools

Sonic Fullbore Imager Dynamic Formation Tester & Corer Magnetic Resonance VSP




MoHoleWS Kanazawa 10/Jun/03/

## Congratulations! Good recovery & good quality Core





MoHoleWS Kanazawa 10/Jun/03/




## Mud Logging / Cuttings Examination

Analysis of the returned drill cuttings gives important and detailed real-time information on the condition of the borehole, and that of the rock being drilled through.











### Sample Time – Depth chart 6000m onland well




出典：石油公団基礎試験報告書

## Low vs. High Fracture Gradient Stress Regimes



MoHoleWS Kanazawa 10/Jun/03/


## Borehole Stability – Operating Mud Window




**ECD: Equivalent Circulation Density**

MoHoleWS Kanazawa 10/Jun/03/

## Real-Time LWD Images for Water-Based Muds



## Borehole Instability – Hole trouble1





MoHoleWS Kanazawa 10/Jun/03/



## Borehole Instability – Hole trouble2



Keep hole straight  
to reduce the risk of  
failing to reach total  
depth




monolews kanazawa 10/Jun/03/

## Consideration for oceanic crust Deep Drilling


- Drilling deep borehole into oceanic crust will require an understanding of the geomechanical processes to ensure **wellbore stability and well control**.
- Operating Mud windows within which to drill deep boreholes will be constrained by **Pore Pressure, Fracture Pressure and Collapse Pressure**.
- **Mud wt windows** : Efficient Hole Cleaning and proper bottom hole Pressure Management are additional operational issues needed to reduce the risk of failing to reach total depth.
- Keep **hole straight** to minimize the torque & drag.
- Drill **a Pilot hole** collecting LWD/PWD, VSP data and Leak off pressure to reduce the risk of failing to reach total depth.
- **APWD**: Real-Time monitoring of borehole and bottom-hole-pressures is highly recommended to ensure wellbore stability and well control.

3/

## Drilling Tools Temperature Limitation



## Riser Operation – NanTroSEIZE Logistics Base



MoHoleWS Kanazawa 10/Jun/03/

## Crew Change by Helicopter (Shorebase - Chikyu)



4 Weeks ON  
4 Weeks OFF

MoHoleWS Kanazawa 10/Jun/03/

## Riser site & Riser hole operation criteria

- **BHST** < 200 - 250°C
- **Water depth** < 4000m - 4500m
- **Weather Window:** 8 months **Low Sea current** and **Calm MetOcean** conditions
- **Seafloor Angle** < 5° for the Riser Wellhead installation
- **Sediment thickness** > 80m - 100m - 36" Conductor jet-in ,  
( **800m** - 20" Casing )
- **Logistics port** < Nearby port **1000km** for Supply Boat
- **Heliport** < Nearby port **350km** for Helicopter Crew Change

MoHoleWS Kanazawa 10/Jun/03/




## Mohole Workshop

# The CHIKYU Engineering Development





Masaoki Yamao  
Engineering Department  
CDEX, JAMSTEC

## “Chikyu” Drilling Records



## NanTroSEIZE Stage 2 (Exp. 319 & 322)

Drilling Location



## Recent Results of Technology Development



## The CHIKYU Enhancement -Targeting to reach Mantle

	Current	Target
Water Depth	2,500m	4,000m+
Total Depth	10,000m	12,000m
Temperature	150 C	300 C
Total Persons	150	200


## Technology Development for Mantle

### Priority Objectives

- Ultra deep hole drilling  
(Target depth : 7,000+mbsf)
- Drilling in ultra deep water  
(Target water depth : 4,000+m)
- High quality core sampling



## Development of Deep Drilling Technology



## 12,000m long Drill Pipes

### Objective :


Development of drill pipes to recover core from upper mantle with 7,000m drilling in 4,000m water

### Results in JFY 2009

- ① Trial production of part of S-155 5-7/8" pipes
- ② Modification of Connection  
Work test with Modified designed thread
- ③ Specimen test with material characteristics using trial production pipe




Trial Production of S-155 5-7/8" Pipes




Trial Piece of Modified Connection

## Measurement Riser Motion



## Next Generation Riser & BOP System



## Technology Roadmap

Area of technologies to be developed		Years required		
		Present ~ 2	+3 ~ 5	+5 ~ 10
Deep Drilling	12 km Drill Pipe	Experimental Model	Manufacturing	Production Model
	Controlled Drilling	Experimental Model	Prototype & MWC <sup>(*)</sup> dev.	Production Model
Deep Water	Riser for Strong Current	Testing & Improvement		
	Deep Water Riser Option:RMR Ssystem	Next Generation Model R&D	Next Generation Experimental Model	

(\*) MWC: Measurement While Coring



*The Mohole Workshop, 3-5 June 2010 @ Kanazawa, Japan*

## ***Technical Development of Coring System in the Quest to reach the Earth's Mantle***


**Yuichi Shinmoto  
Engineering Department  
CDEX JAMSTEC**



### ***Objectives in Technical Development***

- **Enhancement of Core Recovery & Quality**
- **Improvement of Coring Efficiency**
- **Stabilizing & Maintaining Directional Control**
- **Extreme High Temperature Endurance**
- **Prevention of Equipment Fatigue**  
(Drill-pipes, Hydralift Power-Swivel (HPS), etc.)
- **Borehole Stabilization with Mud Circulation**

## Small-Diameter Rotary Core Barrel (SD-RCB)



- Wireline Retrievable
- Core Bit Size (7-1/4"~8-1/2")
- Core Size (3.25") Length (15 ft)
- Diamond Bit Coring in Hard Formation



## Turbo-Corer



Latch-Landing

Motor Section

Bent Housing

Core Tube

Core Bit

### Target Specifications

Description	SI Unit	English Unit
Max. Coring Depth (Below Seafloor)	7000 (m)	23,000 (ft)
Formation Type	Medium ~ Hard	
Core Size	83.0 (mm)	3.25 (in.)
Core Length	4.5 (m)	15 (ft)
Core Bit Size	185 ~ 215 (mm)	7-1/4 ~ 8-1/2 (in.)
Operating WOB	50~100 (kN)	11~22 (k lbs-f)
Max. Operating Torque	7,500 (N-m)	5500 (ft-lbs)
Rotary Speed	100~400 (rpm)	
Max. Temperature	300 (deg.C)	572 (deg.F)
Flow Rate	600 ~ 1,200 (litter/min)	160 ~ 320 (GPM)
Mud Weight	1.0 ~ 1.8 (g/cm <sup>3</sup> )	8.3 ~ 15 (PPG)




## Turbine Performance Test



### Results of Performance Testing

- RPM (800 ~ 8000 rpm)
- Torque (350 N-m) Turbine Only
- Pressure Drop (2MPa)

## Hole Inclinations in NanTroSEIZE Expeditions




- Conventional Drilling BHA: Vertical Hole Inclination varies from 0.70 ~ 4.67 degrees
- Vertical Drilling System (VDS): Hole Inclination of Less than 0.2°
- Hole C0002A: Final Horizontal Displacement of 43 m at the TD



## Development of Directional Control Coring System


*Adjustable Bent Housing middle on the SD-RCB Assembly*



*New 8-1/2" Core Bits*

*Impregnated Diamond & PDC*

## Development Plan of Turbo-Corer and MWC/LWC





# Drilling DEEP into **HOT** Oceanic Crust Constraints and Challenges

David MAINPRICE  
Philippe PEZARD

CNRS, Montpellier (France)




June 3, 2010, Kanazawa (Japan)

## General remarks – stress state of a borehole wall rock

- We are considering horizontal stresses


The stress state of the rock can be divided into 2 components

a) **far field tectonic stress** that classically causes borehole break outs  
(e.g. around the San Andreas fault) – **site dependent** !




b) **local thermo-elastic stresses** caused by changes in the temperature of the wall rock, which lead to brittle failure – **rock dependent** !

## Modeling heat mining for 504B drilling Numerical modeling of the drilling process






Temperature: • Coring:  $\Delta\sigma = \frac{\beta \cdot E}{(1-\nu)} \cdot \Delta T$   
 $\Rightarrow 0.83 \text{ MPa} / ^\circ\text{C} @ 2000 \text{ mbsf}$



- Numerical modeling with LOKI (finite difference)

## Drilling DEEP into HOT Oceanic Crust Constraints & Challenges

- THERMAL
- MECHANICAL
- TECHNOLOGY



- Heat mining process

- **Structure** : heterogeneities (faults, ...), physical properties
- **Dynamics** : pore pressure, stress (**COSOD II**; 1987)

- High Temp (HT) logging tools



## Drilling DEEP into HOT Volcanic Crust What did we learn so far ?

### At sea

ODP (504B) **2111 mbsf** (192°C @ 2004 m)

Hole stopped in a crush zone

Unstable hole from the absence of telescopic casing

No particular problem in the upper crust (normal fault)

Numerous pipe shearing from small active faults

No hole deformation in "normal" drilling conditions

Important hole damages if intensive cooling

Poor core recovery (< 10% at base)



### On land

Kakkonda **450°C**

ISOR < 300°C

IDDP ?

World record in a geothermal context

"Routine" drilling for the past 40 years !


Large difficulties with heterogeneities so far

(and several other locations in the US, Italy, New Zealand, ...)




## Modeling heat mining for 504B drilling Numerical modeling of the drilling process

$$\Delta T = T_{\text{fluid}} - T_{\text{wall rock}}$$



Temperature: • Coring:  $\Delta \sigma = \frac{\beta \cdot E}{(1-\nu)} \cdot \Delta T$   
 $\Rightarrow 0.83 \text{ MPa} / ^\circ\text{C} @ 2000 \text{ mbsf}$




### • Numerical modeling with LOKI (finite difference)




## Modeling heat mining for 504B drilling "Normal" drilling conditions with the JOIDES

Time from the moment the drill pump is started.  $\Delta T = T_{\text{fluid}} - T_{\text{wall rock}}$



Coring Operation - ODP Hole 504B  
Beginning of Leg 148 (Q = 1000 l/min)



Géosciences  
Montpellier  
SUBSURFACE


•  $\Delta T_{\text{max}} = 45^\circ\text{C}$  at 2000 m ( $\Delta S \sim 35$  MPa) : stable hole

## Modeling heat mining for 504B drilling Influence of intense cooling (fishing, ...)


$$\Delta T = T_{\text{fluid}} - T_{\text{wall rock}}$$

collars = base of the pipe

Time from the moment the drill pump is started.



Coring Operation - ODP Hole 504B  
Beginning of Leg 148




• If  $Q = 3400$  liter/min at 2000 m :

$\Delta T_{\text{max}} = 76^\circ\text{C}$  ( $\Delta S \sim 60$  MPa) : tensile failure of the borehole  
(basalt tensile rupture ranges from 50 to 70 MPa after Bauer & Handin, 1985)

Géosciences  
Montpellier  
SUBSURFACE

## Modeling heat mining for 504B drilling Influence of intense cooling (fishing, ...)




Fishing pipe at 1620 m (ODP 140)

Hole cooling at 1980 m (ODP 148)




- Radial, vertical and horizontal thermal ruptures

## Modeling heat mining for 504B drilling Influence of temperature on the coring process



Coring at 2000 m ( $T_{\text{fluid}} \sim 50^\circ\text{C}$  &  $T_{\text{rock}} \sim 190^\circ\text{C}$ ):

the drillbit progresses faster (50 to 100 cm/hr) than the heat ( $\sim 6$  cm/hour) in diabase  $\Rightarrow$  geothermal temperatures are being cored with  $50^\circ\text{C}$  fluid, generating a local temperature gradient of up to  $150^\circ\text{C}$





- $\Delta S > 100$  MPa : the basalt core is shattered at the bit
- the core recovery decreases with depth & temperature







## Heat mining efficiency in the deep oceanic crust Influence of permeability on drilling conditions




- except at shallow depth (and in faults), rock permeability is expected to be low and **drilling conditions undrained**, further weakening the borehole surface

## Heat mining efficiency in the deep oceanic crust Influence of pore pressure and thermal diffusivity




- an **increased pore pressure** will delay the heat penetration into the rock, hence to **weaken the borehole surface**
- the **thermal diffusivity** is expected to decrease with increased temperature, **reinforcing the previous effect**

## Managing heat stress while drilling the oceanic crust Influence of drilling technology in the water column



- **RISER drilling** is the only way to keep the **drilling fluid hot** through the cold ocean ( $2^{\circ}\text{C}$  from 600 m down), hence :
  - to **reduce thermal stresses in the hole**
  - to **increase borehole stability & core recovery**
- **Heat stress THM modelling while drilling** required to adjust the drilling parameters and thereby minimize the thermal stresses in the hole.

## Drilling DEEP into the Oceanic Crust Influence of the structure on borehole stability



- **No major difficulty crossing a 300 m thick normal fault** at shallow depth into the crust (ODP Hole 504B)



## Drilling DEEP into the Oceanic Crust


### Basic HT logging tools requirements for MoHole drilling

- **Temperature surveys :**
  - to measure **in-situ temperature** when re-entering the hole
  - to provide **boundary conditions** for THM numerical modeling
  - to prepare for drilling/logging **operations**
  - to detect the presence of **fluid flow** into/out of the hole
- **Borehole stability control** (from acoustic or "BHTV" images, in order to detect) :
  - **ruptures** of the borehole surface (origine, mapping, ...)
  - identify the **rupture mode** (tensile, compressive ?)
  - detect and map **shear along active fault planes**
  - document changes with depth of the in-situ stress field
  - **plan ahead** drilling operations
- **Temperature and BHTV surveys should be run both at the beginning and at the end of each scientific drilling cruise** to minimize drilling accidents and lost time while at sea.



## Drilling DEEP into the Oceanic Crust

### New HT logging tools : HiTi EC project



Coordinator :  
ISOR (Iceland)

Partners :  
CNRS (France)  
BRGM (France)  
CRES (Greece)  
CALIDUS (UK)  
OXATEC (UK)  
ALT (Lux)  
GFZ (D)

- **HiTi objective** : to plan for **logging as part of IDDP**, to design, built and field test in Iceland **new High Temperature Instruments** for it.
- **Completed & tested so far by HiTi** :
  - **BHTV images** (300°C) and **GR** (300°C) from ALT (wireline)
  - **production tool** (400°C) from CALIDUS (p, T, C<sub>w</sub> tool)
  - **fiber optical monitoring** device (250°C) by GKZ (DTS)
- **To be completed in 2010** :
  - **DLL resistivity** (300°C) from CALIDUS (wireline)
  - **HT temperature** (320°C) from BRGM (wireline)







## • Thermo-elastic stresses: caused by the anisotropy of thermal expansion and elasticity

a) Due to **rapid** changes in wall rock temperature due cooling caused by the circulation of cold drilling fluids (classical phenomenon in Petroleum exploration), near the Moho temperatures are high.

b) Due to **slow** geological cooling, hot asthenosphere to cold lithosphere transition form mid-oceanic ridge axis (e.g. peridotite mantle rock) or cooling from the axial magma chamber (AMC) of a magmatic rock (e.g. oceanic crust composed of basalt or gabbro) – this is general ignored ! But it is responsible for grain boundary cracking.. and can lead to fluid penetration (enhanced cooling, further cracking) and brittle failure.



## Thermo-elastic stresses and grain boundary cracking due to slow geological cooling

The stress intensity factor for tensile mode 1 cracks on a grain boundary (Fredrich and Wong 1986 JGR), depends on;

### **Physical conditions :**

pressure ( $P_c$ ), temperature change  $\Delta T = (T_{\text{plastic}} - T_{\text{wall rock}})$ , is the temperature at which the elastic stress were set to zero.

### **Mineral physical properties :**

For isotropic elasticity (Young's modulus ( $E$ ), Poisson's ratio ( $\nu$ ) and thermal expansion anisotropy

### **Microstructural parameters :**

flaw size ( $a$ ), grain size ( $L = \text{grain size}/2$ )


$$K_{Ic} = \sqrt{\frac{2}{\pi a} \int_0^a \frac{\sigma_{yy}(x) \sqrt{x}}{\sqrt{a-x}} dx} - P_c \sqrt{\pi a}$$

where

$$\sigma_{yy}(x) = \frac{E \Delta T}{2\pi(1-\nu^2)} \left( \frac{4L^2}{4L^2 + (2L-x)^2} - \frac{4L^2}{4L^2 + x^2} + L \ln \left[ \frac{2L-x}{x} \right] - \frac{1}{2} L \ln \left[ \frac{4L^2 (2L-x)^2}{4L^2 + x^2} \right] \right)$$

## Thermo-elastic stresses and grain boundary cracking due to slow geological cooling

Illustration of the method to olivine, predicting more cracking for larger grain size and high temperature differences. Under cooling from  $T_{\text{plastic}} (\approx 900^\circ\text{C}) - T_{\text{wall rock}} (\approx 250^\circ\text{C})$  at Moho gives  $\Delta T = 650^\circ\text{C}$



## Drilling DEEP into **HOT** Oceanic Crust : Constraints & Challenges

### • In-situ thermal stresses :

- can be reduced from **RISER** drilling,
- can have a major and negative impact on **core recovery**,
- will be reduced from **real-time THM modeled** before and during, drilling to define drilling parameters and anticipate failures,

### • Borehole instabilities :

- can be detected and monitored from **repeated BHTV logging**,
- can be reduced from **real-time THM modeling**,
- will be gradually eliminated with **telescopic casing** strings,

### • Site structure and heterogeneities :

- can be detected and monitored from **repeated BHTV logging**,
- have not proven in the past to offer major difficulties in the ocean,

### • HT logging equipment :

- a basic set was developed by HiTI to cover hole stability needs,
- this set requires further tools development to obtain a comprehensive description of the crust at  $T > 200^\circ\text{C}$ .



# ICELAND DEEP DRILLING PROJECT



## What is IDDP?

IDDP is a long-term geothermal research program, established in year 2000


The goal is to improve the economics of geothermal energy by producing supercritical fluids from 4-5 km depth

International Scientific Continental Drilling Program (ICDP) supported meetings and 2 Workshops in 2001 and 2002  
(PI's: GÖF - Wilfred A. Elders and Seiji Seito)

A team of international scientists was established with some 60 research proposals and about 100 - 150 international scientists, engineers and their students from at least 15 countries

**Status 2005 :** Requests for funds for scientific drilling granted by ICDP (1.5 M USD) and by US NSF (3.0 M USD)

**Candidate drillhole:** RN-17 was drilled late in 2004 and completed to 3.1 km depth in February 2005. This well was flow tested in November 2005, and subsequently lost in February 2006 upon cleaning. Cost 4-5 M USD



NOVEMBER 2001- g01



# IDDP Consortium Since 2008




Why are the energy companies interested ?


## Electric Power Generation

	Conventional dry-steam well	IDDP well
Downhole temperature	235 °C	430 - 550 °C
Downhole pressure	30 bar	230 - 260 bar
Volumetric rate of inflow	0.67 m <sup>3</sup> /s	0.67 m <sup>3</sup> /s
Electric power output	~ 5 MW <sub>e</sub>	~ 50 MW <sub>e</sub>

This comparison is based on the same volumetric flow rate of inflowing steam




By drilling deeper we should reach supercritical conditions



# Iceland Deep Drilling Project (IDDP)

- The plan for IDDP is to drill three (3) wells 4.5-5.0 km into the roots of high-temperature geothermal systems in Iceland and seek to reach supercritical conditions  $>374^{\circ}\text{C}$  and  $>22.1\text{ MPa}$ .
- A generic well to achieve the above objectives was designed to serve the dual purpose of a production- and a science well.

## Proposed three IDDP sites





## Production vs. scientific well

Design premises:


- The well should be of sufficient diameter to be suitable as a production well.
- Coring should be done in the expected transition to and inside the supercritical zone, to the extent that science funds would allow.
- Several well designs and coring systems were considered and the conclusion was that the final section below 3500 m should be drilled 8 ½" and 10 m long spot cores collected, rather than the initial idea of continuous coring (HQ) below 2400 m in a 4" diameter hole.

## Expected geological conditions

- The T&P basis of the casing design were boiling point depth (BPD) curve to the Critical Point (~3500 m) and a maximum temp. of 500°C in the well. Lithostatic pressures not expected.
- The wells are to be drilled within three (3) geothermal fields with more than 30 wells each, <3000 m each. Drilling there has been trouble-free, with a few exceptions, and the geological conditions are well known down to 2.5 km.
- The lithology is composed of basaltic lava formations and sub-glacial hyaloclastite formations above ~ 1.5 km depth, and more frequent intrusive rocks, of different type, below that depth.

## T&P design premises

- The static reservoir T&P, the undisturbed conditions at depth.
- Well flowing T&P, dynamic profile.
- Predictions were also made of circulation temperatures for each section of the well at different rig pump flow rates during drilling.




## Conclusions

- The drilling of the IDDP well is a major step for geothermal drilling. Compared to existing HT drilling in Iceland in the past:
  - Extend drilling depth from 3 km to 4.5 km
  - Temperature from 340°C to 400-500°C
  - Cementing casing to twice the depth before.
  - Larger hole and casing sizes.
  - More coring and at extreme temperatures.
  - Four times the cost of a conventional well.

# New challenges

- Risk of intersecting magma and acid fluids, as have been encountered at Krafla over the past few years.
- Greater depths and temperatures than before.
- Circulation temperatures and life of bits.
- Cementing of long casing strings where formation temperatures are high and losses are to be expected.
- Cement design and mud program.
- Casing and connection strength.
- Wellhead temperature and pressure.








## For Coring at Extreme Temperatures, a New Core Barrel was designed and operationally tested for the Iceland Deep Drilling Project

### Expected circulation temperatures of IDDP-1

The return temperatures for 20, 40 and 60 L/s are shown for:

- 2400 m
- 3500 m
- 4500 m



## Design features of the core barrel

- Spot-coring barrel to collect 10 m long cores.
- Diamond impregnated bit 8 ½" od x 4" core.
- Matrix allows for a rotational speed (RPM) between 70 and 160 RPM – actual 70-110 RPM.
- Weight on bit (WOB) should be between 5.4 – 11.4 tonnes – actual 3-10 t.
- Flushing (water) 30 – 40 L/s – actual 25-30 L/s.
- Float valve in core barrel head.
- Temperature logger embedded inside barrel.

## Trial run of IDDP core barrel in RN-17B

- Well RN-17B was in December 2008 being reconditioned by sidetracking it out of well RN-17 at 930 m depth below the production.
- Spot coring was performed at 2800 m depth.
- Rock formation: basalt intrusions.
- Well diameter 12 ¼", inclined 35°.
- Formation temperature is 315°C.
- There were losses near bottom that aided the cooling of the well.

## Týr at well RN-17B on Reykjanes



## Briefing meeting. Crew 6 persons





# Temperature Recording

The Temperature Probe (memory tool) assembled in its housing is attached to the inner core barrel head assembly.

Maximum Temperature Strips made of wax were also inserted into the barrel



The assembled Core Barrel ready for shipping to the Rig site



New spot coring equipment built and tested in 2008 at Reykjanes within in well RN-17B – supported by **ICDP**

New design of core bit – 40 l/s




Six boxes, 1.65 m each, full of 4" drill core recovered from 2500 m depth in RN-17B in 2008






# Measured Depth and ROP




## Weight on bit (WOB) (tonnes) Torque (dNm) and revolution (RPM)





## Drilling Parameters recorded during the Coring Run RN-17B



## Core securely held in Catcher



## Core boxed for examination



## Conclusions


- The barrel performed very well in spite of being deployed in an oversize and inclined hole. Thus slower RPM and less WOB.
- No pilot-hole drilled nor junk-basket run.
- Bit cooling was effective  $<80^{\circ}\text{C}$  – top drive rig.
- Coring rate of penetration 2.6 m/hr.
- 100% core recovery was achieved (9.3 m).
- Overall time for tripping and coring 2 days.

# Drilling Operations at the first Iceland Deep Drilling Well (IDDP)

IDDP #1 Krafla

Saga Rig 0-87 m  
June 2008


- Surface casing
  - 36" hole to 87 m (26" bit and 36" under-reamer)
  - 32" x 0,5" steel casing






# Jötunn Rig 87-260-796 m

Nov-Dec 2008




- Intermediate Casing 1
  - 26 ½" Drillbit
  - 24 ½" Casing 162 lb/ft, K55
- Intermediate Casing 2
  - 23" Drillbit
  - 18 ⅝" Casing 114 lb/ft, K55




# Týr Rig 796 m - final depth

March-July 2009




- 270 ton hook load
- Topdrive
- Automatic pipehandling system
- 3 x Mudpumps
- 180 m<sup>3</sup> mudtanks




## 2009-04-21 Stuck at 2101 m depth

- Pulled out three singles for reaming when hook load dropped suddenly of 20 tons









## 2009-06-02 Coring

- Decided to drill a spot core (9 m) from 2040 m.
- Known loss zones from previous track.
- Managed to drill 2 m until no penetration observed. No core inside the barrel.








## 2009-06-08 Stuck at 2103 m

- Continued drilling until the string got stuck at 2103 m.
- Managed to get loose by pulling 160 tons.
- Circulated for 1 ½" hour.
- RIH again slowly and when the bit was on bottom the torque increased and the string was stuck.
- Returns lost immediately and the string was blocked for circulation.

2009-06-08


Stuck at 2103 m (bit at 2090 m)



2009-06-08

Stuck in Magma at 2104 m


- The Hook load decreases by 50 tons!




## 2009-06-24

# Stuck in Magma!

- At 2100 m the bit was pulled up to casing and the well was cleaned. No fill was on bottom.
- At 2104 m the torque increased and the ROP doubled from 2 m/hr to 4 m/hr.
- The hook load decreased by 40 tons.
- Managed to keep circulation and returns.
- Quenched glass in the returns.
- Proof of Magma.









Krafla, Vítismör  
IDDP-1

16-3-2010  
pEg




Krafla, Vítismör  
IDDP-1


16-3-2010  
pEg



## Future scenario 1:

- Flow test IDDP-1 – as scheduled:
  - 1 Phase 1 test - small scale test (4") **We are here now !**
  - 2 Phase 2 test - full scale test (10")
  - 3 Pilot test for energy production
- Depending on result:
  - 1 Phase 1 successful – go to phase 2 (chemistry and flow OK)
  - 2 Phase 2 successful – go to phase 3 (chemistry and flow OK)
  - 3 Phase 3 successful – produce the well:
    - (i) as a conventional steam producer for the 7 bar inlet pressure to turbine
    - (ii) at higher pressure - into a high-P steam turbine
    - (iii) into a heat exchanger to mitigate chemical problems (cont. pilot study)





The magma chamber(s) in Krafla seen by MT-resistivity survey 2005-2007

## IDDP Plan in 2008

- **IDDP -1** well at Krafla 2008-2009
  - **Landsvirkjun** drills a well to 3.5 km
  - IDDP Consortium deepens the well to 4.5 km
- **IDDP-2** well at Hengill, 2010
  - **Reykjavik Energy** drills a well to 3.5 km
  - IDDP Consortium deepens the well to 5 km
- **IDDP-3** well at Reykjanes, 2011
  - **HS Orka** drills a well to 3.5 km
  - IDDP Consortium deepens the well to 5 km

- Wells at Hengill and Reykjanes are back-up for the IDDP science program

Current Status  
IDDP1 15-20MWe and  
still heating up

Into magma at 2.1 km  
(potential EGS)

Postponed until later

Being considered for  
2011